

PRODUCT INFORMATION

VIEKIRA PAK COMBINATION THERAPY PACK


NAME OF THE MEDICINE

VIEKIRA PAK is a composite pack containing paritaprevir/ritonavir/ombitasvir 75/50/12.5 mg tablets, and dasabuvir 250 mg tablets.

Chemical Structure and Description of each Active Pharmaceutical Ingredient

Paritaprevir

Paritaprevir drug substance is manufactured as a dihydrate, however is dehydrated during the drug product manufacturing process and is amorphous and anhydrous in the final product. Paritaprevir dihydrate is chemically designated (2*R*,6*S*,12*Z*,13*aS*,14*aR*,16*aS*)-*N*-(Cyclopropylsulfonyl)-6-[[[5-methylpyrazin-2-yl)carbonyl]amino]-5,16-dioxo-2-(phenanthridin-6-yloxy)-1,2,3,6,7,8,9,10,11,13*a*,14,15,16,16-tetradecahydrocyclopropa[*e*]pyrrolo[1,2-*a*][1,4]diazacyclopentadecine-14*a*(5*H*)-carboxamide dihydrate. The molecular formula is C₄₀H₄₃N₇O₇S•2H₂O (dihydrate) and the molecular weight for the drug substance is 801.91 (dihydrate). Paritaprevir dihydrate has the following structural formula:




CAS Number: 1456607-71-8

Paritaprevir dihydrate is white to off-white powder with very low water solubility. Paritaprevir dihydrate has pKa of 4.6 at 25°C.

Ritonavir

Ritonavir is chemically designated as [5*S*-(5*R**,8*R**,10*R**,11*R**)]10-Hydroxy-2-methyl-5-(1-methylethyl)-1-[2-(1-methylethyl)-4-thiazolyl]-3,6-dioxo-8,11-bis(phenylmethyl)-2,4,7,12-tetraazatridecan-13-oic acid,5-thiazolylmethyl ester. The molecular formula is C₃₇H₄₈N₆O₅S₂ and the molecular weight is 720.95. Ritonavir has the following structural formula:




CAS Number: 155214-67-5

Ritonavir is a white to off- white to light tan powder practically insoluble in water and freely soluble in methanol and ethanol. Ritonavir has a pKa of 2.8.

Ombitasvir

Ombitasvir drug substance is manufactured as a hydrate, however is dehydrated during the drug product manufacturing process and is amorphous and anhydrous in the final product. Ombitasvir hydrate is chemically designated as Dimethyl (((2*S*,5*S*)-1-(4-*tert*-butylphenyl) pyrrolidine-2,5-diyl)bis{benzene-4,1-diylcarbonyl(2*S*)pyrrolidine-2,1-diyl}[(2*S*)-3-methyl-1-oxobutane-1,2-diyl]})biscarbamate hydrate. The molecular formula is $C_{50}H_{67}N_7O_8 \cdot 4.5H_2O$ (hydrate) and the molecular weight for the drug substance is 975.20 (hydrate). Ombitasvir hydrate has the following structural formula:




CAS Number: 1456607-70-7

Ombitasvir hydrate is white to light pink powder, and is practically insoluble in aqueous buffers but is soluble in ethanol. Ombitasvir hydrate has a pKa of 2.5 at 25°C.

Dasabuvir

Dasabuvir drug substance is manufactured as a sodium salt monohydrate, and is present in the product as the sodium salt monohydrate. Dasabuvir sodium monohydrate is chemically designated as Sodium 3-(3-*tert*-butyl-4-methoxy-5-{6-[(methylsulfonyl)amino]naphthalen-2-yl}phenyl)-2,6-dioxo-3,6-dihydro-2*H*-pyrimidin-1-ide hydrate (1:1:1). The molecular formula is $C_{26}H_{26}N_3O_5S \cdot Na \cdot H_2O$ (salt, hydrate) and the molecular weight of the drug substance is 533.57 (salt, hydrate). Dasabuvir hydrate has the following molecular structure:



CAS Number: 1456607-55-8

Dasabuvir sodium monohydrate is white to off-white to pink powder, slightly soluble in water and very slightly soluble in methanol and isopropyl alcohol. The pKa values of dasabuvir are 8.2 (pK₁) and 9.2 (pK₂).

DESCRIPTION

Paritaprevir, ritonavir, and ombitasvir are co-formulated as film-coated immediate release tablets. The tablets also contain copovidone, tocopherol, propylene glycol monolaurate, sorbitan monolaurate, silicon dioxide, sodium stearyl fumarate and Opadry II pink 85F140088 (polyvinyl alcohol, titanium dioxide, macrogol, purified talc, and iron oxide red). The tablets do not contain gluten. The strength for the fixed dose combination tablet is 75 mg paritaprevir/50 mg ritonavir/12.5 mg ombitasvir.

Dasabuvir is formulated as a 250 mg film-coated, immediate release tablet containing microcrystalline cellulose, lactose, copovidone, croscarmellose sodium, colloidal anhydrous silica, magnesium stearate, and Opadry II Beige 85F97497 (polyvinyl alcohol, titanium dioxide, macrogol, purified talc, and iron oxide yellow, iron oxide red and iron oxide black.). The tablets do not contain gluten.

PHARMACOLOGY

Pharmacodynamics

Pharmacotherapeutic group: Direct-acting antiviral, ATC code: J05AX66

Mechanism of Action

VIEKIRA PAK combines three direct-acting hepatitis C virus antiviral agents with distinct mechanisms of action and non-overlapping resistance profiles to target HCV at multiple steps in the viral lifecycle.

Paritaprevir

Paritaprevir is an inhibitor of HCV NS3/4A protease which is necessary for the proteolytic cleavage of the HCV encoded polyproteins (into mature forms of the NS3, NS4A, NS4B, NS5A, and NS5B proteins) and is essential for viral replication.

Ritonavir is not active against HCV. Ritonavir is a pharmacokinetic enhancer that increases peak and trough plasma drug concentrations of paritaprevir and overall drug exposure (i.e. area under the curve).

Ombitasvir

Ombitasvir is an inhibitor of HCV NS5A which is necessary for viral replication.

Dasabuvir

Dasabuvir is a non-nucleoside inhibitor of the HCV RNA-dependent RNA polymerase encoded by the NS5B gene.

Activity in Cell Culture and/or Biochemical Studies

Paritaprevir

In a biochemical assay, paritaprevir inhibited the proteolytic activity of the recombinant HCV genotype 1a and 1b NS3/4A protease enzymes with IC_{50} values of 0.18 nM and 0.43 nM, respectively. The EC_{50} of paritaprevir against genotype 1a-H77 and 1b-Con1 strains in the HCV replicon cell culture assay was 1.0 and 0.21 nM, respectively. The activity of paritaprevir was attenuated 24- to 27-fold in the presence of 40% human plasma. The mean EC_{50} of paritaprevir against replicons containing NS3 from a panel of treatment-naïve genotype 1a and 1b isolates in the HCV replicon cell culture assay was 0.86 nM (range 0.43 to 1.87 nM; $n = 11$) and 0.06 nM (range 0.03 to 0.09 nM; $n = 9$), respectively. Paritaprevir had an EC_{50} value of 5.3 nM against the 2a-JFH-1 replicon cell line, and EC_{50} values of 19, 0.09, and 0.68 nM against replicon cell lines containing NS3 from a single isolate each of genotype 3a, 4a, and 6a, respectively. In a biochemical assay, paritaprevir inhibited the activity of NS3/4A enzymes from single isolates of genotypes 2a, 2b, 3a, and 4a with IC_{50} values of 2.4, 6.3, 14.5, and 0.16 nM, respectively.

Ritonavir did not exhibit a direct antiviral effect on the replication of HCV subgenomic replicons, and the presence of ritonavir did not affect the *in vitro* antiviral activity of paritaprevir.

Ombitasvir

In replicon cell culture assays, ombitasvir has EC_{50} values of 14.1 pM and 5.0 pM against HCV genotypes 1a-H77 and 1b-Con1, respectively. The activity of ombitasvir was attenuated 11- to 13-fold in the presence of 40% human plasma. The mean EC_{50} of ombitasvir against replicons containing NS5A from a panel of treatment-naïve genotype 1a and 1b isolates in the HCV replicon cell culture assay was 0.66 pM (range 0.35 to 0.88 pM; $n = 11$) and 1.0 pM (range 0.74 to 1.5 pM; $n = 11$), respectively. Ombitasvir has EC_{50} values of 12, 4.3, 19, 1.7, 3.2, and 366 pM against replicon cell lines constructed with NS5A from single isolates representing genotypes 2a, 2b, 3a, 4a, 5a, and 6a, respectively. Negligible anti-viral activity against genotypes 1a-H77 and 1b-Con1 was noted by the human major metabolites of ombitasvir, M29 and M36 in the HCV replicon assay; M29 and M36 do not contribute to antiviral activity of ombitasvir.

Dasabuvir

In a biochemical assay, dasabuvir inhibited the polymerase activity of the recombinant HCV genotype 1a and 1b HCV NS5B enzymes with IC_{50} values of 2.8 nM and 10.7 nM, respectively. The EC_{50} of dasabuvir against genotype 1a-H77 and 1b-Con1 strains in HCV replicon cell culture assays was 7.7 and 1.8 nM, respectively. The replicon activity of dasabuvir was attenuated 12- to 13-fold in the presence of 40% human plasma. The mean EC_{50} of dasabuvir against replicons containing NS5B from a panel of treatment-naïve genotype 1a and 1b isolates in the HCV replicon cell culture assay was 0.77 nM (range 0.4 to 2.1 nM; $n = 11$) and 0.46 nM (range 0.2 to 2 nM; $n = 10$), respectively. In biochemical assays, dasabuvir inhibited a panel of genotype 1a and 1b polymerases with a mean IC_{50} value of 4.2 nM (range 2.2 to 10.7 nM; $n = 7$). Dasabuvir had lower potency (>200 times) against polymerases from other HCV genotypes (2a, 2b, 3a and 4a). The M1 metabolite of dasabuvir had 30–40% lower potency than dasabuvir against genotypes 1a-H77 and 1b-Con1 in the HCV replicon assay.

Combination Activity *in vitro*

All two-drug combinations of paritaprevir, ombitasvir, dasabuvir and ribavirin demonstrated additive to synergistic inhibition of HCV genotype 1 replicon at the majority of drug concentrations studied in short term cell culture assays. In long term replicon survival assays, the ability of drug-resistant cells to form colonies in the presence of a single drug or drugs in combination was evaluated. In pair-wise combinations of paritaprevir, ombitasvir, and dasabuvir at concentrations 10-fold over their respective EC₅₀, colony survival was reduced by more than 100-fold by two drugs as compared to each drug alone. When all three drugs were combined at concentrations of 5-fold above their respective EC₅₀, no drug-resistant colonies survived.

Resistance in Cell Culture

Resistance to paritaprevir, ombitasvir, or dasabuvir conferred by variants in NS3, NS5A, or NS5B, respectively, selected in cell culture or identified in Phase 2b and 3 clinical trials were phenotypically characterized in the appropriate genotype 1a or 1b replicons.

In genotype 1a, substitutions F43L, R155 G/K/S, A156T, and D168A/E/F/H/N/V/Y in HCV NS3 reduced susceptibility to paritaprevir by 7- to 219-fold. The activity of paritaprevir in genotype 1a was not significantly affected (less than or equal to 3-fold) by single substitutions V23A (in NS4A), V36A/M, V55I, Y56H, Q80K or E357K. Double variants including combinations of V36M, F43L, Y56H, or E357K with R155K or with a D168 substitution reduced the activity of paritaprevir by an additional 2- to 7-fold relative to the single R155K or D168 substitution. In genotype 1b, substitutions A156T, D168A/H/V/Y, and Y56H in combination with D168A/V/Y in HCV NS3 reduced susceptibility to paritaprevir. In the genotype 1b replicon, the activity of paritaprevir was reduced 27- to 337-fold by D168A/H/V/Y substitutions. The combination of Y56H and D168A, D168V or D168Y reduced the activity of paritaprevir by an additional 12- to 26-fold relative to the single D168 substitution in genotype 1b replicons.

In genotype 1a, substitutions M28T/V, Q30E/R, H58D, Y93C/H/L/N in HCV NS5A reduced susceptibility to ombitasvir by 58- to 67,000 fold. In genotype 1b, substitutions L28T, L31F/V, and Y93H in HCV NS5A reduced susceptibility to ombitasvir 8- to 661 fold. In general, combinations of ombitasvir resistance-associated substitutions in HCV genotype 1a or 1b replicons further reduced ombitasvir antiviral activity.

In genotype 1a, substitutions C316Y, M414I/T, N444K, E446K/Q, Y448C/H, A553T, G554S, S556G/R, and Y561H in HCV NS5B reduced susceptibility to dasabuvir by 5- to 1472 fold. G558R and D559G/N were observed as treatment-emergent substitutions but the activity of dasabuvir against these variants could not be evaluated due to poor replication capacity. In genotype 1b, substitutions C316H/N/Y, S368T, N411S, M414I/T/V, Y448C/H, A553V, S556G and D559G in HCV NS5B reduced susceptibility to dasabuvir by 5- to 1569 fold. Dasabuvir retained full activity against replicons containing substitutions S282T in the nucleoside binding site, M423T in the lower thumb site, and P495A/S, P496S or V499A in the upper thumb site.

Effect of Baseline HCV Substitutions/Polymorphisms on Treatment Response

A pooled analysis of subjects in the Phase 2b and 3 clinical trials treated with paritaprevir, ombitasvir, and dasabuvir with or without ribavirin was conducted to explore the association between the baseline NS3/4A, NS5A or NS5B substitutions/polymorphisms and treatment outcome in recommended regimens.

In the greater than 500 genotype 1a baseline samples in this analysis, the most frequently observed resistance-associated variants were M28V (7.4%) in NS5A and S556G (2.9%) in NS5B. Q80K, although

a highly prevalent polymorphism in NS3 (41.2% of samples), confers minimal resistance to paritaprevir. Resistance-associated variants at amino acid positions R155 and D168 in NS3 were rarely observed (less than 1%) at baseline. In the greater than 200 genotype 1b baseline samples in this analysis, the most frequently observed resistance-associated variants observed were Y93H (7.5%) in NS5A, and C316N (17.0%) and S556G (15%) in NS5B. Given the low virologic failure rates observed with recommended treatment regimens for HCV genotype 1a- and 1b-infected subjects, the presence of baseline variants appears to have little impact on the likelihood of achieving SVR.

Resistance in Clinical Studies

Of the 2,510 HCV genotype 1 infected subjects in the Phase 2b and 3 clinical trials treated with regimens containing paritaprevir, ombitasvir, and dasabuvir with or without ribavirin (for 8, 12, or 24 weeks), a total of 74 subjects (3%) experienced virologic failure (primarily post-treatment relapse). Treatment-emergent variants and their prevalence in these virologic failure populations are shown in Table 1. In the 67 genotype 1a infected subjects, NS3 variants were observed in 50 subjects, NS5A variants were observed in 46 subjects, NS5B variants were observed in 37 subjects, and treatment-emergent variants were seen in all 3 drug targets in 30 subjects. In the 7 genotype 1b infected subjects, treatment-emergent variants were observed in NS3 in 4 subjects, in NS5A in 2 subjects, and in both NS3 and NS5A in 1 subject. No genotype 1b infected subjects had treatment-emergent variants in all 3 drug targets.

Table 1. Treatment-Emergent Amino Acid Substitutions in the Pooled Analysis of VIEKIRA PAK with and without Ribavirin Regimens in Phase 2b and Phase 3 Clinical Trials (N = 2510)

Target	Emergent Amino Acid Substitutions ^a	Genotype 1a N = 67 ^b % (n)	Genotype 1b N = 7 % (n)
NS3	V55I ^c	6 (4)	-
	Y56H ^c	9 (6)	42.9 (3) ^d
	I132V ^c	6 (4)	-
	R155K	13.4 (9)	-
	D168A	6 (4)	-
	D168V	50.7 (34)	42.9 (3) ^d
	D168Y	7.5 (5)	-
	V36A ^c , V36M ^c , F43L ^c , D168H, E357K ^c	< 5%	-
NS5A	M28T	20.9 (14)	-
	M28V ^e	9 (6)	-
	Q30R ^e	40.3 (27)	-
	Y93H	-	28.6 (2)
	H58D, H58P, Y93N	< 5%	-
NS5B	A553T	6.1 (4)	-
	S556G	33.3 (22)	-
	C316Y, M414T, G554S, S556R, G558R, D559G, D559N, Y561H	< 5%	-

a. Observed in at least 2 subjects of the same subtype.

b. N = 66 for the NS5B target.

c. Substitutions were observed in combination with other emergent substitutions at NS3 position R155 or D168.

d. Observed in combination in genotype 1b-infected subjects.

e. Observed in combination in 6% (4/67) of the subjects.

Note: The following variants were selected in cell culture but were not treatment-emergent: NS3 variants A156T in genotype 1a, and R155Q and D168H in genotype 1b; NS5A variants Y93C/H in genotype 1a, and L31F/V or Y93H in combination with L28M, L31F/V or P58S in genotype 1b; and NS5B variants Y448H in

Attachment 1: Product information AusPAR VIEKIRA PAK - Paritaprevir / Ritonavir / Ombitasvir and Dasabuvir (as Sodium Salt) - AbbVie Pty Ltd - PM-2014-01436-1-2 FINAL 24 March 2017. This Product Information was approved at the time this AusPAR was published.

Target	Emergent Amino Acid Substitutions ^a	Genotype 1a N = 67 ^b % (n)	Genotype 1b N = 7 % (n)
	genotype 1a, and M414T and Y448H in genotype 1b.		

Persistence of Resistance-Associated Substitutions

The persistence of paritaprevir, ombitasvir, and dasabuvir resistance-associated amino acid substitutions in NS3, NS5A, and NS5B, respectively, was assessed in genotype 1a-infected subjects in Phase 2b trials. Paritaprevir treatment-emergent variants V36A/M, R155K or D168V were observed in NS3 in 47 subjects. Ombitasvir treatment-emergent variants M28T, M28V or Q30R in NS5A were observed in 32 subjects. Dasabuvir treatment-emergent variants M414T, G554S, S556G, G558R or D559G/N in NS5B were observed in 34 subjects.

NS3 variants V36A/M and R155K and NS5B variants M414T and S556G remained detectable at post-treatment Week 48, whereas NS3 variant D168V and all other NS5B variants were not observed at post-treatment Week 48. All treatment-emergent variants in NS5A remained detectable at post-treatment Week 48. Due to high SVR rates in genotype 1b, trends in persistence of treatment-emergent variants in this genotype could not be established.

The lack of detection of virus containing a resistance-associated substitution does not indicate that the resistant virus is no longer present at clinically significant levels. The long-term clinical impact of the emergence or persistence of virus containing VIEKIRA PAK -resistance-associated substitutions is unknown.

Cross-resistance

Cross-resistance is expected among NS5A inhibitors, NS3/4A protease inhibitors, and non-nucleoside NS5B inhibitors by class. The impact of prior ombitasvir, paritaprevir or dasabuvir treatment experience on the efficacy of other NS5A inhibitors, NS3/4A protease inhibitors, or NS5B inhibitors has not been studied.

Pharmacodynamic interactions

Coadministration with enzyme inducers may increase the risk of adverse events and ALT elevations. Coadministration with ethinylestradiol may increase the risk of ALT elevations (see INTERACTIONS WITH OTHER MEDICINES).

Pharmacokinetics

The pharmacokinetic properties of the combination of paritaprevir, ombitasvir, ritonavir, and dasabuvir have been evaluated in healthy adult subjects and in subjects with chronic hepatitis C. Table 2 shows mean C_{max} and AUC_{0-24} of paritaprevir/ritonavir/ombitasvir 150/100/25 mg once daily with dasabuvir 250 mg twice daily following multiple doses with food in healthy volunteers.

Table 2: Geometric Mean C_{max} and AUC_{0-24} of Multiple Doses of paritaprevir/ritonavir/ombitasvir 150/100/25 mg Once Daily with dasabuvir 250 mg Twice Daily with Food in Healthy Volunteers

	C_{max} (ng/mL)	AUC_{0-24} (ng*hr/mL)
paritaprevir	1470	6990

ombitasvir	127	1420
dasabuvir	1030	13680
ritonavir	1600	9470

Absorption

Paritaprevir/ritonavir/ombitasvir and dasabuvir were absorbed after oral administration with mean T_{max} of approximately 4 to 5 hours. While ombitasvir and dasabuvir exposures increased in a dose proportional manner, paritaprevir and ritonavir exposures increased in a more than dose proportional manner. Accumulation is minimal for ombitasvir and dasabuvir and approximately 1.5- to 2-fold for ritonavir and paritaprevir. Pharmacokinetic steady state for the combination is achieved after approximately 12 days of dosing.

Effects of Food on Oral Absorption

Paritaprevir, ritonavir, ombitasvir and dasabuvir should be administered with food. All clinical trials with paritaprevir, ritonavir, ombitasvir and dasabuvir have been conducted following administration with food.

Food increased the exposure (AUC) of paritaprevir, ombitasvir, ritonavir, and dasabuvir by up to 211%, 82%, 49%, and 30% respectively relative to the fasting state. The increase in exposure was similar regardless of meal type (e.g., high-fat versus moderate-fat) or calorie content (approximately 600 Kcal versus approximately 1000 Kcal). To maximise absorption, VIEKIRA PAK should be taken with food without regard to fat or calorie content.

Distribution

Paritaprevir, ombitasvir, ritonavir and dasabuvir are highly bound to plasma proteins. Plasma protein binding is not meaningfully altered in patients with renal or hepatic impairment. The blood to plasma concentration ratios in humans ranged from 0.5 to 0.7, indicating that paritaprevir, ombitasvir, and dasabuvir were preferentially distributed in the plasma compartment of whole blood. Paritaprevir was approximately 97 to 98.6% bound to human plasma proteins over a concentration range of 0.08 to 8 microgram/mL. Ritonavir was greater than 99% bound to human plasma proteins over a concentration range of 0.07-22 microgram/mL. Ombitasvir was approximately 99.9% bound to human plasma proteins over a concentration range of 0.09 to 9 microgram/mL. Dasabuvir was > 99.5% bound to human plasma proteins over a concentration range of 0.15 to 5 microgram/mL.

In animals, paritaprevir liver levels are significantly higher than plasma levels (e.g. liver: plasma ratio of >300:1 in mouse). *In vitro* data indicate that paritaprevir is a substrate for the human hepatic uptake transporters, OATP1B1 and OATP1B3.

Metabolism

Paritaprevir

Paritaprevir is metabolised predominantly by CYP3A4 and to a lesser extent CYP3A5. Following administration of a single 200/100 mg oral dose of ^{14}C paritaprevir/ritonavir to humans, the parent drug was the major circulating component accounting for approximately 90% of the plasma radioactivity. At least 5 minor metabolites of paritaprevir have been identified in circulation that

accounted for approximately 10% of plasma radioactivity. These metabolites are not expected to have antiviral activity.

Ombitasvir

Ombitasvir is metabolised via amide hydrolysis followed by oxidative metabolism. Following a 25 mg single dose of ¹⁴C-ombitasvir given alone, unchanged parent drug accounted for 8.9% of total radioactivity in human plasma; a total of 13 metabolites were identified in human plasma. These metabolites are not expected to have antiviral activity or off-target pharmacological activity.

Dasabuvir

Dasabuvir is predominantly metabolised by CYP2C8 and to a lesser extent by CYP3A. Following a 400 mg ¹⁴C-dasabuvir dose in humans, unchanged dasabuvir was the major component (approximately 60%) of drug related radioactivity in plasma; seven metabolites were identified in plasma. The most abundant plasma metabolite was M1, which represented 21% of drug-related radioactivity (AUC) in circulation and has similar contribution to activity against genotype 1 as the parent drug after accounting for difference in protein binding.

Ritonavir

Ritonavir is predominantly metabolised by CYP3A and to a lesser extent, by CYP2D6. Nearly the entire plasma radioactivity after a single 600 mg dose of ¹⁴C-ritonavir oral solution in humans was attributed to unchanged ritonavir.

Elimination

Paritaprevir

Following dosing of paritaprevir/ritonavir/ombitasvir with or without dasabuvir, mean plasma half-life of paritaprevir was approximately 5.5 hours. Following a 200 mg ¹⁴C-paritaprevir dose with 100 mg ritonavir, approximately 88% of the radioactivity was recovered in faeces with limited radioactivity (8.8%) in urine.

Ombitasvir

Following dosing of paritaprevir/ritonavir/ombitasvir with or without dasabuvir, mean plasma half-life of ombitasvir was approximately 21-25 hours. Following a 25 mg ¹⁴C-ombitasvir dose, approximately 90.2% of the radioactivity was recovered in faeces with limited radioactivity (1.91%) in urine.

Dasabuvir

Following dosing of dasabuvir with paritaprevir/ritonavir/ombitasvir, mean plasma half-life of dasabuvir was approximately 5.5 to 6 hours. Following a 400 mg ¹⁴C-dasabuvir dose, approximately 94.4% of the radioactivity was recovered in faeces with limited radioactivity (approximately 2%) in urine.

Ritonavir

Following dosing of paritaprevir/ritonavir/ombitasvir, mean plasma half-life of ritonavir was approximately 4 hours. Following a 600 mg dose of ¹⁴C-ritonavir oral solution, 86.4% of the radioactivity was recovered in the faeces and 11.3% of the dose was excreted in the urine.

Implications for Drug Interactions

Potential for VIEKIRA PAK to affect the pharmacokinetics of other medicinal products

In vivo drug interaction studies evaluated the net effect of the combination treatment, including ritonavir.

The following section describes the specific transporters and metabolizing enzymes that are affected by VIEKIRA PAK. See INTERACTIONS WITH OTHER MEDICINES for guidance regarding potential interactions with other medicinal products and dosing recommendations.

Medicinal products metabolised by CYP3A4

Ritonavir is a strong inhibitor of CYP3A. Co-administration of VIEKIRA PAK with medicinal products primarily metabolised by CYP3A may result in increased plasma concentrations of these medicinal products. Medicinal products that are highly dependent on CYP3A for clearance and for which elevated plasma levels are associated with serious events are contraindicated (see CONTRAINDICATIONS and INTERACTIONS WITH OTHER MEDICINES).

CYP3A substrates evaluated in drug interaction studies which may require dose adjustment and/or clinical monitoring include (see Table 13) cyclosporine, tacrolimus, amlodipine, rilpivirine and alprazolam. Examples of other CYP3A4 substrates which may require dose adjustment and/or clinical monitoring include calcium channel blockers (e.g. nifedipine), and trazodone. Although buprenorphine and zolpidem are also metabolised by CYP3A, drug interaction studies indicate that no dose adjustment is needed when co-administering these medicinal products with VIEKIRA PAK (see Table 13).

Medicinal products transported by the OATP family and OCT1

Paritaprevir is an inhibitor of the hepatic uptake transporters OATP1B1 and OATP1B3, and paritaprevir and ritonavir are inhibitors of OATP2B1. Ritonavir is an *in vitro* inhibitor of OCT1, but the clinical relevance is unknown. Co-administration of VIEKIRA PAK with medicinal products that are substrates of OATP1B1, OATP1B3, OATP2B1 or OCT1 may increase plasma concentrations of these transporter substrates, potentially requiring dose adjustment/clinical monitoring. Such medicinal products include some statins (see Table 13), fexofenadine, repaglinide and angiotensin II receptor antagonists (e.g., valsartan).

OATP1B1/3 substrates evaluated in drug interaction studies include pravastatin and rosuvastatin (see Table 13).

Medicinal products transported by BCRP

Paritaprevir, ritonavir and dasabuvir are inhibitors of BCRP *in vivo*. Co-administration of VIEKIRA PAK together with medicinal products that are substrates of BCRP may increase plasma concentrations of these transporter substrates, potentially requiring dose adjustment/clinical monitoring. Such medicinal products include sulfasalazine, imatinib and some of the statins (see Table 13).

BCRP substrates evaluated in drug interaction studies include rosuvastatin (see Table 13).

Medicinal products transported by P-gp in the intestine

While paritaprevir, ritonavir and dasabuvir are *in vitro* inhibitors of P-gp, no significant change was observed in the exposure of the P-gp substrate digoxin when administered with VIEKIRA PAK and dasabuvir (see Table 13). VIEKIRA PAK may increase the plasma exposure to medicinal products that are sensitive for changed intestinal P-gp activity (such as dabigatran etexilate).

Medicinal products metabolised by glucuronidation (UGT1A1)

Paritaprevir, ombitasvir and dasabuvir are inhibitors of UGT1A1. Co-administration of VIEKIRA PAK with medicinal products that are primarily metabolised by UGT1A1 result in increased plasma concentrations of such medicinal products; routine clinical monitoring is recommended for narrow therapeutic index medicinal products (i.e. levothyroxine). See also INTERACTIONS WITH OTHER MEDICINES for specific advice on raltegravir and buprenorphine, which have been evaluated in drug interaction studies.

Medicinal products metabolised by CYP2C19

Co-administration of VIEKIRA PAK can decrease exposures of medicinal products that are metabolised by CYP2C19 (e.g. lansoprazole, esomeprazole, s-mephenytoin), which may require dose adjustment/clinical monitoring. CYP2C19 substrates evaluated in drug interaction studies include omeprazole and escitalopram (see Table 13).

Medicinal products metabolised by CYP2C9

VIEKIRA PAK did not affect the exposures of the CYP2C9 substrate, warfarin. Other CYP2C9 substrates (NSAIDs (e.g. ibuprofen), antidiabetics (e.g. glimepiride, glipizide) are not expected to require dose adjustments.

Medicinal products metabolised by CYP2D6 or CYP1A2

VIEKIRA PAK did not affect the exposures of the CYP2D6/CYP1A2 substrate, duloxetine. Other CYP1A2 substrates (e.g. ciprofloxacin, theophylline and caffeine) and CYP2D6 substrates (e.g. desipramine, metoprolol and dextromethorphan) are not expected to require dose adjustments.

Medicinal products renally excreted via transport proteins

Ombitasvir, paritaprevir, and ritonavir do not inhibit organic anion transporter (OAT1) *in vivo* as shown by the lack of interaction with tenofovir (OAT1 substrate). *In vitro* studies show that ombitasvir, paritaprevir, and ritonavir are not inhibitors of organic cation transporters (OCT2), organic anion transporters (OAT3), or multidrug and toxin extrusion proteins (MATE1 and MATE2K) at clinically relevant concentrations.

Therefore, VIEKIRA PAK is not expected to affect medicinal products which are primarily excreted by the renal route via these transporters.

Potential for other medicinal products to affect the pharmacokinetics of ombitasvir, paritaprevir, and dasabuvir

Medicinal products that inhibit CYP3A4

Co-administration of VIEKIRA PAK with strong inhibitors of CYP3A may increase paritaprevir concentrations (see Table 13).

Enzyme inducers

Co-administration of VIEKIRA PAK with medicinal products that are moderate or strong enzyme inducers is expected to decrease ombitasvir, paritaprevir, ritonavir and dasabuvir plasma

concentrations and reduce their therapeutic effect. Contraindicated enzyme inducers are provided under CONTRAINDICATIONS and INTERACTIONS WITH OTHER MEDICINES.

Medicinal products that inhibit CYP3A4 and transport proteins

Paritaprevir is eliminated via CYP3A4 mediated metabolism and biliary excretion (substrate of the hepatic transporters OATP1B1, P-gp and BCRP). Caution is advised if co-administering VIEKIRA PAK with medicinal products that are both moderate inhibitors of CYP3A4 and inhibitors of multiple transporters (P-gp, BCRP and/or OATP1B1/ OATP1B3). These medicinal products may show clinically relevant increases in exposures of paritaprevir (e.g., ritonavir with atazanavir, erythromycin, diltiazem or verapamil).

Medicinal products that inhibit transport proteins

Potent inhibitors of P-gp, BCRP, OATP1B1 and/or OATP1B3 have the potential to increase the exposure to paritaprevir. Inhibition of these transporters is not expected to show clinically relevant increases in exposures of ombitasvir and dasabuvir.

Special Populations

Renal Impairment

Paritaprevir/ritonavir/ombitasvir and dasabuvir

Based on the pharmacokinetic data in HCV uninfected subjects (n=24), no dose adjustment of VIEKIRA PAK is recommended in subjects with mild, moderate or severe renal impairment. The efficacy and safety of VIEKIRA PAK have not been evaluated in HCV-infected subjects with moderate or severe renal impairment. Pharmacokinetics of the combination of paritaprevir 150 mg, ombitasvir 25 mg, and ritonavir 100 mg, with or without dasabuvir 400 mg were evaluated in subjects with mild (CrCl: 60 to 89 mL/min), moderate (CrCl: 30 to 59 mL/min) and severe (CrCl: 15 to 29 mL/min) renal impairment.

In subjects with mild renal impairment (n=6), paritaprevir mean C_{max} and AUC values were comparable (up to 19% higher), ombitasvir mean C_{max} and AUC values were comparable (up to 7% lower), and ritonavir mean C_{max} and AUC values were 26% to 42% higher and dasabuvir mean C_{max} and AUC values were 5% to 21% higher compared to subjects with normal renal function.

In subjects with moderate renal impairment (n=6), paritaprevir mean C_{max} values were comparable (< 1% increase) and AUC values were 33% higher, ombitasvir mean C_{max} and AUC values were comparable (up to 12% lower), and ritonavir mean C_{max} and AUC value were 48% to 80% and dasabuvir mean C_{max} and AUC values were 9% to 37% higher compared to subjects with normal renal function.

In subjects with severe renal impairment (n=6), paritaprevir mean C_{max} values were comparable (< 1% increase) and AUC values were 45% higher, ombitasvir mean C_{max} and AUC values were comparable (up to 15% lower), and ritonavir mean C_{max} and AUC value were 66% to 114% higher and dasabuvir mean C_{max} and AUC values were 12% to 50% higher compared to subjects with normal renal function.

Hepatic Impairment

The changes in paritaprevir, ombitasvir, dasabuvir and ritonavir exposures in subjects with mild and moderate hepatic impairment are not considered clinically significant. No dose adjustment for VIEKIRA PAK is recommended in HCV-infected subjects with mild and moderate hepatic impairment. VIEKIRA PAK is contraindicated in patients with severe hepatic impairment.

Pharmacokinetics of the combination of paritaprevir 200 mg, and ritonavir 100 mg, ombitasvir 25 mg, and dasabuvir 400 mg were evaluated in subjects (n=17) with mild (Child-Pugh A), moderate (Child-Pugh B) and severe (Child-Pugh C) hepatic impairment.

In subjects with mild hepatic impairment (n=6), paritaprevir, ritonavir and ombitasvir mean C_{max} and AUC values decreased by 29% to 48%, 34% to 40% and up to 8%, respectively, and dasabuvir mean C_{max} and AUC values were 17% to 24% higher compared to subjects with normal hepatic function.

In subjects with moderate hepatic impairment (n=6), paritaprevir mean C_{max} and AUC value increased by 26% to 62%, ombitasvir and ritonavir mean C_{max} and AUC values decreased by 29% to 30% and 30 to 33%, respectively, and dasabuvir mean C_{max} and AUC values were 16% to 39% lower compared to subjects with normal hepatic function. The safety and efficacy of VIEKIRA PAK have not been established in HCV-infected patients with moderate hepatic impairment (Child-Pugh B); however, no dose adjustment is expected to be required based on pharmacokinetic studies.

In subjects with severe hepatic impairment (n=5), paritaprevir and dasabuvir mean C_{max} and AUC values increased by 3.2 to 9.5-fold and 0.3- to 3.3-fold respectively; ritonavir mean C_{max} values were 35% lower and AUC values were 13% higher and ombitasvir mean C_{max} and AUC values decreased by 68% and 54% respectively compared to subjects with normal hepatic function.

Elderly

No dose adjustment is necessary for paritaprevir/ritonavir/ombitasvir or dasabuvir in elderly patients.

Based on population pharmacokinetic analysis of data from Phase 3 clinical studies, a 10 year increase or decrease in age from 54 years (median age in the Phase 3 studies) would result in approximately 10% change in ombitasvir exposures, less than 10% change in dasabuvir exposures and $\leq 20\%$ change in paritaprevir exposures. There is no pharmacokinetic information in patients >75 years.

Paediatric Population (<18 years)

The pharmacokinetics, safety and efficacy of VIEKIRA PAK in paediatric patients have not been established.

Race or Ethnicity

No dose adjustment is necessary for paritaprevir/ritonavir/ombitasvir or dasabuvir based on race or ethnicity. Based on population pharmacokinetic analysis of data from Phase 3 clinical studies, Asian subjects had 18% to 21%, 37% to 39% and 29% to 39% higher ombitasvir, paritaprevir and dasabuvir exposures, respectively, than non-Asian subjects. The ritonavir exposures were comparable between Asians and non-Asians. However, patient numbers in the clinical trials were not sufficient to definitively address possible differences in pharmacokinetics and toxicity profiles in specific ethnic groups such as Asian patients.

Sex or Body weight

No dose adjustment is necessary for paritaprevir/ritonavir/ombitasvir or dasabuvir based on gender or body weight.

Based on population pharmacokinetic analysis of data from Phase 3 clinical studies, female subjects would have approximately 55%, 100%, 15% and 21% higher ombitasvir, paritaprevir, ritonavir and dasabuvir exposures (AUC), respectively, than male subjects. A 10 kg change in body weight from

76 kg (median weight in the Phase 3 studies) would results in <10% change in ombitasvir and dasabuvir exposures, and no change in paritaprevir exposures. Body weight is not a significant predictor of ritonavir exposures.

CLINICAL TRIALS

The efficacy and safety of VIEKIRA PAK were evaluated in six randomised Phase 3 clinical trials, in over 2,300 subjects with genotype 1 chronic hepatitis C infection. Included in the Phase 3 program was one trial exclusively in subjects with cirrhosis (Child-Pugh A). Phase 3 trials are summarised in Table 3.

Table 3: Phase 3 Randomised, Global Multicentre Trials Conducted with VIEKIRA PAK with or without ribavirin (RBV).

Trial¹	Number of subjects (treated²)	HCV Genotype (GT)	Summary of Study Design³
Treatment-naïve⁴, without cirrhosis			
SAPPHIRE I	631	GT1	Arm A: VIEKIRA PAK + RBV Arm B: Placebo
PEARL III	419	GT1b	Arm A: VIEKIRA PAK+ RBV Arm B: VIEKIRA PAK
PEARL IV	305	GT1a	Arm A: VIEKIRA PAK+ RBV Arm B: VIEKIRA PAK
Treatment-experienced⁵, without cirrhosis			
SAPPHIRE II	394	GT1	Arm A: VIEKIRA PAK + RBV Arm B: Placebo
PEARL II (open-label)	180	GT1b	Arm A: VIEKIRA PAK+ RBV Arm B: VIEKIRA PAK
Treatment-naïve and treatment-experienced⁵, with compensated cirrhosis			
TURQUOISE II (open-label)	380	GT1	Arm A: VIEKIRA PAK + RBV (12 weeks) Arm B: VIEKIRA PAK + RBV (24 weeks)
1 Double-blind unless otherwise noted			
2 Treated is defined as subjects who were randomised and received at least one dose of VIEKIRA PAK.			
3 Treatment duration was 12 weeks for all arms, except for TURQUOISE II which included a 24 week arm.			
4 Treatment naïve was defined as not having received any prior therapy for HCV infection.			
5 Treatment-experienced subjects were defined as either: prior relapsers (subjects with HCV RNA undetectable at or after the end of at least 36 weeks of pegIFN/RBV treatment, but HCV RNA was detectable within 52 weeks of treatment follow-up) or prior partial responders (received at least 20 weeks of pegIFN/RBV and achieved a greater than or equal to 2 log ₁₀ IU/mL reduction in HCV RNA at week 12, but not achieving HCV RNA undetectable at end of treatment) or prior null-responders (received at least 12 weeks of pegIFN/RBV treatment and failed to achieve a 2 log ₁₀ IU/mL reduction in HCV RNA at week 12 or received at least 4 weeks of pegIFN/RBV treatment and achieved a < 1 log ₁₀ IU/mL reduction in HCV RNA at week 4).			

In all six trials, the paritaprevir/ritonavir/ombitasvir dose was 150/100/25 mg once daily and the dasabuvir dose was 250 mg twice daily. For subjects who received ribavirin, the ribavirin dose was 1000 mg per day for subjects weighing less than 75 kg or 1200 mg per day for subjects weighing greater than or equal to 75 kg.

Sustained virologic response (virologic cure) was defined as unquantifiable or undetectable HCV RNA 12 weeks after the end of treatment (SVR12) in the Phase 3 trials. Treatment duration was fixed in each trial and was not guided by subjects' HCV RNA levels (no response-guided algorithm). Plasma HCV RNA values were measured during the clinical trials using the COBAS TaqMan HCV test (version 2.0), for use with the High Pure System. The assay had a lower limit of quantification (LLOQ) of 25 IU per mL.

Pooled Analyses of Clinical Trials

Durability of Response

Overall, 660 subjects in Phase 2 and 3 clinical trials had HCV RNA results for both the SVR12 and SVR24 time points. Among these subjects, the positive predictive value of SVR12 on SVR24 was 99.8%.

Pooled Efficacy Analyses

In phase 3 clinical trials, 1096 subjects (including 202 with cirrhosis) received the recommended regimen for their HCV subtype, cirrhosis status and previous treatment. Table 4 shows SVR rates for these patients.

Among subjects who received the recommended regimen, 97% achieved SVR (95% with cirrhosis and 97% without cirrhosis), while 0.5% demonstrated virologic breakthrough and 1.6% experienced post-treatment relapse.

Table 4: SVR12 rates for recommended treatment regimens

	Genotype 1a		Genotype 1b	
	No Cirrhosis VIEKIRA PAK- RBV	With Cirrhosis VIEKIRA PAK-RBV	No cirrhosis VIEKIRA PAK	With cirrhosis VIEKIRA PAK-RBV
	12 weeks	12 weeks*	12 weeks	12 weeks
Treatment-naïve	96% (403/420)	92% (61/66)	100% (210/210)	100% (22/22)
Treatment-experienced	96% (166/173)	94% (64/68)*	100% (91/91)	98% (45/46)
Prior pegIFN/RBV relapser	94% (47/50)	93% (14/15)	100% (33/33)	100% (14/14)
Prior pegIFN/RBV partial responder	100% (36/36)	100% (11/11)	100% (26/26)	86% (6/7)
Prior pegIFN/RBV null responder	95% (83/87)	93% (39/42) (24 weeks)	100% (32/32)	100% (25/25)
TOTAL	96% (569/593)	93% (125/134)*	100% (301/301)	99% (67/68)
*All subjects received 12 weeks of therapy except for genotype 1a infected prior null responders with cirrhosis who received 24 weeks of therapy.				

Impact of Ribavirin Dose Adjustment on Probability of SVR

In Phase 3 clinical trials, 91.5% of subjects did not require ribavirin dose adjustments during therapy. In the 8.5% of subjects who had ribavirin dose adjustments during therapy, the SVR rate (98.5%) was comparable to subjects who maintained their starting ribavirin dose throughout treatment.

Clinical Trials in Treatment-Naïve Adults

SAPPHIRE-I (M11-646) - Genotype 1, Treatment-Naïve

SAPPHIRE-I was a randomised, global multicentre, double-blind, placebo-controlled trial conducted in 631 treatment-naïve adults with genotype 1 chronic hepatitis C virus infection without cirrhosis. VIEKIRA PAK was given for 12 weeks of treatment. Subjects randomised to the placebo arm received placebo for 12 weeks, after which they received open-label VIEKIRA PAK with ribavirin for 12 weeks.

Treated subjects (N=631) had a median age of 52 years (range: 18 to 70); 64.8% were born between 1945 – 1965; 54.5% were male; 5.4% were Black and 5.1% were Hispanic or Latino; 16.2% had a body mass index of at least 30 kg/m²; 15.2% had a history of depression or bipolar disorder; 69.3% had IL28B non-CC genotype; 79.1% had baseline HCV RNA levels at least 800,000 IU/mL; 15.4% had portal fibrosis (F2) and 8.7% had bridging fibrosis (F3); 67.7% had HCV genotype 1a infection; 32.3% had HCV genotype 1b infection.

Table 5 shows the SVR12 rates for genotype 1-infected, treatment-naïve subjects receiving VIEKIRA PAK-RBV for 12 weeks in SAPPHIRE-I.

Table 5: SVR12 for Genotype 1-Infected Treatment-Naïve Subjects in SAPPHIRE-I

Treatment Outcome	VIEKIRA PAK –RBV for 12 Weeks		
	n/N	%	95% CI
Overall SVR12	456/473	96.4	94.7, 98.1
HCV genotype 1a	308/322	95.7	93.4, 97.9
HCV genotype 1b	148/151	98.0	95.8, 100.0
Outcome for subjects without SVR12			
On-treatment VF ^a	1/473	0.2	
Relapse ^b	7/463	1.5	
Other ^c	9/473	1.9	
CI = confidence interval, VF = virologic failure			
a. On-treatment VF was defined as confirmed HCV ≥ 25 IU/mL after HCV RNA < 25 IU/mL during treatment, confirmed 1 log ₁₀ IU/mL increase in HCV RNA from nadir, or HCV RNA persistently ≥ 25 IU/mL with at least 6 weeks of treatment.			
b. Relapse was defined as confirmed HCV RNA ≥ 25 IU/mL post-treatment before or during SVR12 window among subjects with HCV RNA < 25 IU/mL at last observation during at least 11 weeks of treatment.			
c. Other includes subjects not achieving SVR12 but not experiencing on-treatment VF or relapse (e.g. missing HCV RNA values in the SVR12 window).			

In the primary efficacy analysis, VIEKIRA PAK-RBV demonstrated superiority to the historical control SVR rate of 78% {95% CI of 75%, 80%} (based upon telaprevir plus peginterferon (pegIFN/RBV) for

subjects with genotype 1 HCV infection who were treatment-naïve without cirrhosis. Refer to the telaprevir prescribing information.

No subjects with HCV genotype 1b infection experienced on-treatment virologic failure and one subject with HCV genotype 1b infection experienced relapse.

Based on subgroup analyses, the following baseline factors were not associated with lower SVR12 rates (lower 95% confidence bound > 70%):

- *Viral factors:* genotype 1 subtype, baseline viral load
- *Host factors:* Gender, race, ethnicity, age, birth year (1945 – 1965), IL28B allele, baseline body mass index, history of depression or bipolar disorder, fibrosis stage

In addition, subjects who underwent ribavirin dose modifications did not have lower SVR12 rates.

Significantly more subjects (352/363 = 97.0%) who received VIEKIRA PAK with ribavirin had normalised ALT by the end of treatment than those who received placebo (18/114 = 15.8%); *P* value < 0.001.

PEARL-III (M13-961) – Genotype 1b, Treatment-Naïve

PEARL-III was a randomised, global multi-centre, double-blind, controlled trial conducted in 419 treatment-naïve adults with genotype 1b chronic hepatitis C virus infection without cirrhosis. Subjects were randomised in a 1:1 ratio to receive VIEKIRA PAK or VIEKIRA-PAK-RBV for 12 weeks of treatment.

Treated subjects (N=419) had a median age of 50 years (range: 19 to 70); 54.9% were born between 1945 – 1965, 45.8% were male; 4.8% were Black; 1.7% were Hispanic or Latino; 16.5% had a body mass index of at least 30 kg/m²; 9.3% had a history of depression or bipolar disorder; 79.0% had IL28B non-CC genotype; 73.3% had baseline HCV RNA of at least 800,000 IU/mL; 20.3% had portal fibrosis (F2) and 10.0% had bridging fibrosis (F3).

Table 6 shows the SVR12 rates for genotype 1b-infected, treatment-naïve subjects who received VIEKIRA PAK or VIEKIRA PAK-RBV for 12 weeks in PEARL III. In this study, VIEKIRA PAK had similar SVR12 rates (100%) compared to VIEKIRA PAK with ribavirin (99.5%).

Table 6: SVR12 for Genotype 1b-Infected Treatment-Naïve Subjects in PEARL III

	VIEKIRA PAK-RBV for 12 weeks			VIEKIRA PAK for 12 weeks		
	n/N	%	95% CI	n/N	%	95% CI
Overall SVR12	209/210	99.5	98.6, 100.0	209/209	100	98.2, 100.0
Outcome for subjects without SVR12	1/210	0.5		2/209	1.0	
On-treatment VF ^a	1/210	0.5		0/209	0	
Relapse ^b	0/210	0		0/209	0	
Other ^c	0/210	0		0/209	0	

CI = confidence interval, VF = virologic failure

- On-treatment VF was defined as confirmed HCV \geq 25 IU/mL after HCV RNA $<$ 25 IU/mL during treatment, confirmed 1 log₁₀ IU/mL increase in HCV RNA from nadir, or HCV RNA persistently \geq 25 IU/mL with at least 6 weeks of treatment.
- Relapse was defined as confirmed HCV RNA \geq 25 IU/mL post-treatment before or during SVR12 window among subjects with HCV RNA $<$ 25 IU/mL at last observation during at least 11 weeks of treatment.
- Other includes subjects not achieving SVR12 but not experiencing on-treatment VF or relapse (e.g. missing HCV RNA values in the SVR12 window).

In the primary efficacy analysis, VIEKIRA PAK and VIEKIRA PAK with ribavirin demonstrated superiority to the historical control SVR rate of 80% {95% CI of 75%, 84%} (based upon telaprevir plus pegIFN/RBV) for subjects with genotype 1b HCV infection who were treatment-naïve without cirrhosis. Refer to the telaprevir prescribing information.

Based on subgroup analyses, the following baseline factors were not associated with lower SVR12 rates (lower 95% confidence bound $>$ 73%):

- Viral factors:* baseline viral load
- Host factors:* Gender, race, ethnicity, age, birth year (1945 – 1965), IL28B allele, baseline body mass index, history of depression or bipolar disorder, fibrosis stage

In addition, subjects who underwent ribavirin dose modifications did not have lower SVR12 rates.

PEARL-IV (M14-002) – Genotype 1a, Treatment-Naïve

PEARL-IV was a randomised, global multicentre, double-blind, controlled trial conducted in 305 treatment-naïve adults with genotype 1a chronic hepatitis C virus infection without cirrhosis. Subjects were randomised in a 1:2 ratio to receive VIEKIRA PAK or VIEKIRA PAK with ribavirin for 12 weeks of treatment.

Treated subjects (N=305) had a median age of 54 years (range: 19 to 70); 72.5% were born between 1945 – 1965, 65.2% were male; 11.8% were Black; 9.2% were Hispanic or Latino; 19.7% had a body mass index of at least 30 kg/m²; 20.7% had a history of depression or bipolar disorder; 69.2% had IL28B non-CC genotype; 86.6% had baseline HCV RNA levels of at least 800,000 IU/mL; 18.4% had portal fibrosis (F2) and 17.7% had bridging fibrosis (F3).

Table 7 shows the SVR12 rates for genotype 1a-infected, treatment-naïve subjects who received VIEKIRA PAK or VIEKIRA PAK-RBV for 12 weeks in PEARL IV. VIEKIRA PAK was not non-inferior to VIEKIRA PAK with ribavirin.

Table 7: SVR12 for Genotype 1a-Infected Treatment-Naïve Subjects in PEARL IV

	VIEKIRA PAK RBV for 12 weeks			VIEKIRA PAK for 12 weeks		
	n/N	%	95% CI	n/N	%	95% CI
Overall SVR12	97/100	97.0	93.7, 100.0	185/205	90.2	86.2, 94.3
Outcome for subjects without SVR12						
On-treatment VF ^a	1/100	1.0		6/205	2.9	
Relapse ^b	1/98	1.0		10/194	5.2	
Other ^c	1/100	1.0		1/205	0.5	

CI = confidence interval, VF = virologic failure

- a. On-treatment VF was defined as confirmed HCV \geq 25 IU/mL after HCV RNA $<$ 25 IU/mL during treatment, confirmed 1 log₁₀ IU/mL increase in HCV RNA from nadir, or HCV RNA persistently \geq 25 IU/mL with at least 6 weeks of treatment.
- b. Relapse was defined as confirmed HCV RNA \geq 25 IU/mL post-treatment before or during SVR12 window among subjects with HCV RNA $<$ 25 IU/mL at last observation during at least 11 weeks of treatment.
- c. Other includes subjects not achieving SVR12 but not experiencing on-treatment VF or relapse (e.g. missing HCV RNA values in the SVR12 window).

In the primary efficacy analysis, VIEKIRA PAK and VIEKIRA PAK with ribavirin demonstrated superiority to the historical control SVR rate of 72% {95% CI of 68%, 75%} (based upon telaprevir plus pegIFN/RBV) for subjects with genotype 1a HCV infection who were treatment-naïve without cirrhosis. Refer to the telaprevir prescribing information.

Based on subgroup analyses, the following baseline factors were not associated with lower SVR12 rates (lower 95% confidence bound $>$ 65%) :

- *Viral factors*: baseline viral load
- *Host factors*: Gender, race, ethnicity, age, birth year (1945 – 1965), IL28B allele, baseline body mass index, history of depression or bipolar disorder, fibrosis stage

In addition, subjects who underwent ribavirin dose modifications did not have lower SVR12 rates.

Clinical Trials in Treatment-Experienced Adults

SAPPHIRE-II (M13-098) Genotype 1 – Treatment-experienced

SAPPHIRE-II was a randomised, global multicentre, double-blind, placebo-controlled trial conducted in 394 subjects with genotype 1 chronic hepatitis C virus infection without cirrhosis who did not achieve SVR with prior treatment with pegIFN/RBV. VIEKIRA PAK-RBV was given for 12 weeks of treatment. Subjects randomised to the placebo arm received placebo for 12 weeks, after which they received VIEKIRA PAK-RBV for 12 weeks.

Treated subjects (N=394) had a median age of 54 years (range: 19 to 71); 49.0% were prior pegIFN/RBV null responders; 21.8% were prior pegIFN/RBV partial responders, and 29.2% were prior pegIFN/RBV relapsers; 73.9% were born between 1945 – 1965; 57.6% were male; 8.1% were Black and 6.3% were Hispanic or Latino; 19.8% had a body mass index of at least 30 kg/m²; 20.6% had a history of depression or bipolar disorder; 89.6% had IL28B non-CC genotype; 87.1% had baseline HCV RNA levels at least 800,000 IU per mL; 17.8% had portal fibrosis (F2) and 14.5% had bridging fibrosis (F3); 58.4% had HCV genotype 1a infection; 41.4% had HCV genotype 1b infection.

Table 8 shows the SVR12 rates for treatment-experienced subjects with genotype 1-infection receiving VIEKIRA PAK with ribavirin for 12 weeks in SAPPHIRE-II.

Table 8: SVR12 for Genotype 1-infected Treatment-Experienced Subjects in SAPPHERE-II

Treatment Outcome	VIEKIRA PAK -RBVfor 12 weeks		
	n/N	%	95% CI
Overall SVR12	286/297	96.3	94.1, 98.4
HCV Genotype 1a	166/173	96.0	93.0, 98.9
Prior pegIFN/RBV null responder	83/87	95.4	91.0, 99.8
Prior pegIFN/RBV partial responder	36/36	100	100.0, 100.0
Prior pegIFN/RBV relapser	47/50	94.0	87.4, 100.0
HCV Genotype 1b	119/123	96.7	93.6, 99.9
Prior pegIFN/RBV null responder	56/59	94.9	89.3, 100.0
Prior pegIFN/RBV partial responder	28/28	100	100.0, 100.0
Prior pegIFN/RBV relapser	35/36	97.2	91.9, 100.0
Outcome for subjects without SVR12			
On-treatment VF ^a	0/297	0	
Relapse ^b	7/293	2.4	
Other ^c	4/297	1.3	

CI = confidence interval, VF = virologic failure

a. On-treatment VF was defined as confirmed HCV \geq 25 IU/mL after HCV RNA < 25 IU/mL during treatment, confirmed 1 log₁₀ IU/mL increase in HCV RNA from nadir, or HCV RNA persistently \geq 25 IU/mL with at least 6 weeks of treatment.

b. Relapse was defined as confirmed HCV RNA \geq 25 IU/mL post-treatment before or during SVR12 window among subjects with HCV RNA < 25 IU/mL at last observation during at least 11 weeks of treatment.

c. Other includes subjects not achieving SVR12 but not experiencing on-treatment VF or relapse (e.g. missing HCV RNA values in the SVR12 window).

No subjects with HCV genotype 1b infection experienced on-treatment virologic failure and 2 subjects with HCV genotype 1b infection experienced relapse.

In the primary efficacy analysis, VIEKIRA PAK with ribavirin demonstrated superiority to the historical control SVR rate of 65% {95% CI of 60%, 70%} (based upon telaprevir plus pegIFN/RBV) for subjects with genotype 1 HCV infection who were treatment-experienced without cirrhosis. Refer to the telaprevir prescribing information.

Significantly more subjects (217/224 = 96.9%) who received VIEKIRA PAK with ribavirin had normalised ALT by the end of treatment than those who received placebo (Arm B, 10/78=12.8%); *P* value < 0.001.

Based on subgroup analyses, the following baseline factors were not associated with lower SVR12 rates (lower 95% confidence bound > 60%):

- *Viral factors*: genotype 1 subtype, baseline viral load
- *Host factors*: prior pegIFN/RBV response, gender, race, ethnicity, age, birth year (1945 – 1965), IL28B allele, baseline body mass index, history of depression or bipolar disorder, fibrosis stage

In addition, subjects who underwent ribavirin dose modifications did not have lower SVR12 rates.

PEARL-II (M13-389) – Genotype 1b, Treatment-Experienced

PEARL-II was a randomised, global multicentre, open-label trial conducted in 180 adults with chronic genotype 1b hepatitis C virus infection without cirrhosis who did not achieve SVR with prior treatment with pegIFN/RBV. Subjects were randomised, in a 1:1 ratio, to receive VIEKIRA PAK or VIEKIRA PAK-RBV for 12 weeks of treatment.

Treated subjects (N=179) had a median age of 57 years (range: 26 to 70); 35.2% were prior pegIFN/RBV null responders; 28.5% were prior pegIFN/RBV partial responders, and 36.3% were prior pegIFN/RBV relapsers; 70.9% were born between 1945 – 1965; 54.2% were male; 3.9% were Black; 1.7% were Hispanic or Latino; 21.8% had a body mass index of at least 30 kg/m²; 12.8% had a history of depression or bipolar disorder; 90.5% had IL28B non-CC genotype; 87.7% had baseline HCV RNA levels of at least 800,000 IU/mL; 17.9% had portal fibrosis (F2) and 14.0% had bridging fibrosis (F3).

Table 9 shows the SVR12 rates for genotype 1b-infected, treatment-experienced subjects who received VIEKIRA PAK or VIEKIRA PAK-RBV for 12 weeks in PEARL II. In this study, VIEKIRA PAK had a similar SVR12 rate (100%) compared to VIEKIRA PAK with ribavirin (97.7%).

Table 9: SVR12 for Genotype 1b-infected Treatment-Experienced Subjects in PEARL II

	VIEKIRA PAK-RBV for 12 weeks			VIEKIRAPAK for 12 weeks		
	n/N	%	95% CI	n/N	%	95% CI
Overall SVR12	86/88	97.7	94.6, 100.0	91/91	100	95.9, 100.0
Prior pegIFN/RBV null responder	30/31	96.8	90.6, 100.0	32/32	100	89.3, 100.0
Prior pegIFN/RBV partial responder	24/25	96.0	88.3, 100.0	26/26	100	87.1, 100.0
Prior pegIFN/RBV relapser	32/32	100	89.3, 100.0	33/33	100	89.6, 100.0
Outcome for subjects without SVR12						
On-treatment VF ^a	0/88	0		0/91	0	
Relapse ^b	0/88	0		0/91	0	
Other ^c	2/88	2.3		0/91	0	

CI = confidence interval, VF = virologic failure

- On-treatment VF was defined as confirmed HCV ≥ 25 IU/mL after HCV RNA < 25 IU/mL during treatment, confirmed 1 log₁₀ IU/mL increase in HCV RNA from nadir, or HCV RNA persistently ≥ 25 IU/mL with at least 6 weeks of treatment.
- Relapse was defined as confirmed HCV RNA greater than 25 IU/mL post-treatment before or during SVR12 window among subjects with HCV RNA < 25 IU/mL at last observation during at least 11 weeks of treatment.
- Other includes subjects not achieving SVR12 but not experiencing on-treatment VF or relapse (e.g. missing HCV RNA values in the SVR12 window).

In the primary efficacy analysis, VIEKIRA PAK and VIEKIRA PAK with ribavirin demonstrated superiority to the historical control SVR rate of 69% {95% CI of 62%, 75%} (based upon telaprevir plus

pegIFN/RBV) for subjects with genotype 1b HCV infection who were treatment- experienced without cirrhosis. Refer to the telaprevir prescribing information.

Based on subgroup analyses, the following baseline factors were not associated with lower SVR12 rates (lower 95% confidence bound > 64%):

- *Viral factors:* baseline viral load
- *Host factors:* prior pegIFN/RBV response, gender, race, ethnicity, age, birth year (1945 – 1965), IL28B allele, baseline body mass index, history of depression or bipolar disorder, fibrosis stage

In addition, subjects who underwent ribavirin dose modifications did not have lower SVR12 rates.

Clinical Trial in Subjects with Cirrhosis

TURQUOISE-II (M13-099) – Genotype 1, Treatment-naïve or treatment-experienced patients with cirrhosis

TURQUOISE-II was a randomised, global multicentre, open-label trial conducted exclusively in 380 genotype 1-infected subjects with cirrhosis (Child-Pugh A) who were either treatment-naïve or did not achieve SVR with prior treatment with pegIFN/RBV. VIEKIRA PAK with ribavirin was administered for either 12 or 24 weeks of treatment.

Treated subjects (N=380) had a median age of 58 years (range: 21 to 71); 42.1% were treatment-naïve, 36.1% were prior pegIFN/RBV null responders; 8.2% were prior pegIFN/RBV partial responders, 13.7% were prior pegIFN/RBV relapsers; 85.5% were born between 1945 – 1965; 70.3% were male; 3.2% were Black; 11.8% were Hispanic or Latino; 28.4% had a body mass index of at least 30 kg/m²; 14.7% had platelet counts of < 90 x 10⁹/L; 11.3% had albumin (< 35 g/L); 86.1% had baseline HCV RNA levels of at least 800,000 IU/mL; 81.8% had IL28B non-CC genotype; 24.7% had a history of depression or bipolar disorder; 68.7% had HCV genotype 1a infection, 31.3% had HCV genotype 1b infection.

Table 10 shows the SVR12 rates for genotype 1-infected subjects with cirrhosis who were treatment-naïve or previously treated with pegIFN/RBV.

Table 10: SVR12 for Genotype 1-Infected Subjects with Cirrhosis who were Treatment-Naïve or Previously Treated with pegIFN/RBV

Treatment Outcome	VIEKIRA PAK -RBV					
	12 Weeks			24 Weeks		
	n/N	%	CI ^a	n/N	%	CI ^a
Overall SVR12	191/208	91.8	87.6, 96.1	166/172	96.5	93.4, 99.6
HCV Genotype 1a	124/140	88.6	83.3, 93.8	115/121	95.0	91.2, 98.9
Treatment naïve	59/64	92.2		53/56	94.6	
Prior pegIFN/RBV null responders	40/50	80.0		39/42	92.9	
Prior pegIFN/RBV partial responders	11/11	100		10/10	100	
Prior pegIFN/RBV Prior relapsers	14/15	93.3		13/13	100	
HCV Genotype 1b	67/68	98.5	95.7, 100	51/51	100	93.0, 100
Treatment naïve	22/22	100		18/18	100	
Prior pegIFN/RBV null responders	25/25	100		20/20	100	
Prior pegIFN/RBV partial responders	6/7	85.7		3/3	100	
Prior pegIFN/RBV Prior relapsers	14/14	100		10/10	100	
Outcome for subjects without SVR12						
On-treatment VF ^b	1/208	0.5		3/172	1.7	
Relapse ^c	12/203	5.9		1/164	0.6	
Other ^d	4/208	1.9		2/172	1.2	

CI = confidence interval, VF = virologic failure, NA = data not yet available

- 97.5% confidence intervals are used for the primary efficacy endpoints (overall SVR12 rate); 95% confidence intervals are used for additional efficacy endpoints (SVR12 rates in HCV genotype 1a and 1b subjects).
- On-treatment VF was defined as confirmed HCV \geq 25 IU/mL after HCV RNA < 25 IU/mL during treatment, confirmed 1 log₁₀ IU/mL increase in HCV RNA from nadir, or HCV RNA persistently \geq 25 IU/mL with at least 6 weeks of treatment.
- Relapse was defined as confirmed HCV RNA \geq 25 IU/mL post-treatment before or during SVR12 window among subjects with HCV RNA < 25 IU/mL at last observation during at least 11 or 22 weeks of treatment, for subjects assigned to 12 or 24 weeks of treatment, respectively.
- Other includes subjects not achieving SVR12 but not experiencing on-treatment VF or relapse (e.g. missing HCV RNA values in the SVR12 window).

In the primary efficacy analysis, VIEKIRA PAK-RBV administered for 12 or 24 weeks demonstrated superiority to the historical control SVR rate of 47% {95% CI of 41%, 54%} (based upon telaprevir plus pegIFN/RBV) for subjects with genotype 1 HCV infection with cirrhosis that were treatment-naïve or previously treated with pegIFN/RBV. Refer to the telaprevir prescribing information.

Based on subgroup analyses, the following baseline factors were not associated with lower SVR12 rates (lower 95% confidence bound > 43%):

- *Viral factors:* genotype 1 subtype, baseline viral load
- *Host factors:* prior pegIFN/RBV response, gender, ethnicity, age, birth year (1945 – 1965), IL28B allele, baseline body mass index, history of depression or bipolar disorder, fibrosis stage, baseline platelet count, baseline albumin

In addition, subjects who underwent ribavirin dose modifications did not have lower SVR12 rates.

INDICATIONS

VIEKIRA PAK is indicated for the treatment of genotype 1 chronic hepatitis C infection, including patients with compensated cirrhosis. Duration of therapy and addition of ribavirin are dependent on patient population (see DOSAGE AND ADMINISTRATION, PRECAUTIONS, CLINICAL TRIALS).

CONTRAINDICATIONS

Hypersensitivity to components of VIEKIRA PAK, or to any of the excipients.

Patients with severe hepatic impairment (Child-Pugh C).

Use of ethinylestradiol-containing medicinal products such as those contained in most combined oral contraceptives or contraceptive vaginal rings

Drugs that are highly dependent on CYP3A for clearance and for which elevated plasma concentrations are associated with serious and/or life-threatening events.

Drugs that are moderate or strong inducers of CYP3A and strong inducers of CYP2C8 and may lead to reduced efficacy of VIEKIRA PAK.

Drugs that are strong inhibitors of CYP2C8 and may increase dasabuvir plasma concentrations and the risk of QT prolongations.

The following drugs are contraindicated with VIEKIRA PAK.

Table 11 Drugs that are Contraindicated with VIEKIRA PAK

Drug Class	Drug(s) within Class that are Contraindicated	Clinical Comments
Alpha1- adrenoceptor antagonist	Alfuzosin HCL	Potential for hypotension.
Antiarrhythmics	Amiodarone, quinidine	Potential for cardiac arrhythmias.
Anticonvulsants	Carbamazepine, phenytoin, phenobarbital	Ombitasvir, paritaprevir, ritonavir and dasabuvir exposures may decrease leading to a potential loss of therapeutic activity of VIEKIRA PAK.
Antihistamines (for systemic use)	Astemizole, terfenadine	Potential for cardiac arrhythmias.
GI motility agent	Cisapride	Potential for cardiac arrhythmias.

Attachment 1: Product information AusPAR VIEKIRA PAK - Paritaprevir / Ritonavir / Ombitasvir and Dasabuvir (as Sodium Salt) - AbbVie Pty Ltd - PM-2014-01436-1-2 FINAL 24 March 2017. This Product Information was approved at the time this AusPAR was published.

Antigout medications	Colchicine (in patients with renal or hepatic impairment)	Increased potential for colchicine associated adverse events.
Antihyperlipidemic agent	Gemfibrozil	Increase in dasabuvir exposures by 10-fold which may increase the risk of QT prolongation.
Antimycobacterial	Rifampicin	Ombitasvir, paritaprevir, ritonavir and dasabuvir exposures may decrease leading to a potential loss of therapeutic activity of VIEKIRA PAK.
Antipsychotic	Blonanserin	No information on potential effects is currently available
Ergot derivatives	Ergotamine, dihydroergotamine, ergonovine, methylergonovine	Acute ergot toxicity characterized by vasospasm and tissue ischemia has been associated with co-administration of ritonavir and ergonovine, ergotamine, dihydroergotamine, or methylergonovine.
Ethinyl estradiol-containing products	Ethinyl estradiol-containing medications such as combined oral contraceptives	Potential for ALT elevations (see PRECAUTIONS).
Herbal Product	St. John's Wort (<i>Hypericum perforatum</i>)	Ombitasvir, paritaprevir, ritonavir and dasabuvir exposures may decrease leading to a potential loss of therapeutic activity of VIEKIRA PAK.
HMG-CoA Reductase Inhibitors	Lovastatin, simvastatin	Potential for myopathy including rhabdomyolysis.
Long Acting Beta-Adrenoceptor agonist	Salmeterol	The combination may result in increased risk of cardiovascular adverse events associated with salmeterol, including QT prolongation, palpitations and sinus tachycardia.
Neuroleptics	Pimozide	Potential for cardiac arrhythmias.
Non-nucleoside reverse transcriptase inhibitor	Efavirenz	Co-administration of efavirenz based regimens with paritaprevir, ritonavir plus dasabuvir was poorly tolerated and resulted in liver enzyme elevations.
Phosphodiesterase-5 (PDE5) inhibitor	Sildenafil when dosed for the treatment of pulmonary arterial hypertension (PAH)	There is increased potential for sildenafil-associated adverse events such as visual disturbances, hypotension, priapism, and syncope.
Platelet aggregation inhibitors excluding heparin	Ticagrelor	Increased potential for ticagrelor associated adverse events.

Sedatives/hypnotics	Triazolam Orally administered midazolam	Triazolam and orally administered midazolam are extensively metabolised by CYP3A4. Coadministration of triazolam or orally administered midazolam with VIEKIRA PAK may cause large increases in the concentration of these benzodiazepines. The potential exists for serious and/or life threatening events such as prolonged or increased sedation or respiratory depression.
Steroid antibacterials	Fusidic acid	Increased potential for fusidic acid associated adverse events.
Anticancer agents	Mitotane Enzalutamide	Increased potential for mitotane and enzalutamide associated adverse events.

PRECAUTIONS

VIEKIRA PAK efficacy has not been studied in patients who have previously failed therapy with a treatment regimen that includes VIEKIRA PAK or other direct-acting antiviral agents.

ALT Elevations

During clinical trials with VIEKIRA PAK with or without ribavirin, transient, asymptomatic elevations of alanine transaminase (ALT) to greater than 5 times the upper limit of normal (ULN) occurred in approximately 1% of all subjects (see ADVERSE REACTIONS). These ALT elevations were significantly more frequent in female subjects who were using ethinyl estradiol-containing medications such as combined oral contraceptives, contraceptive patches, or contraceptive vaginal rings (see CONTRAINDICATIONS). ALT elevations typically occurred during the first 4 weeks of treatment and declined within approximately two weeks of onset with continued dosing of VIEKIRA PAK with or without ribavirin.

Ethinyl estradiol-containing medications must be discontinued prior to starting therapy with VIEKIRA PAK (see CONTRAINDICATIONS). Alternative contraceptive agents or methods of contraception (e.g., progestin only contraception or non-hormonal methods) are recommended during VIEKIRA PAK therapy. Ethinyl estradiol-containing medications can be restarted approximately 2 weeks following completion of treatment with VIEKIRA PAK.

Subjects using estrogens other than ethinyl estradiol, such as estradiol and conjugated estrogens used in hormone replacement therapy had a rate of ALT elevation similar to those not receiving any estrogens (1%). No additional monitoring of ALT is required outside of local recommendations and routine clinical practice guidelines.

If ALT is found to be elevated above baseline levels, it should be monitored closely.

- Patients should be instructed to consult their health care professional without delay if they have onset of fatigue, weakness, lack of appetite, nausea and vomiting, jaundice or discoloured urine or faeces (see Serum Bilirubin Elevations under ADVERSE REACTIONS).

- Discontinue VIEKIRA PAK if ALT elevation is accompanied by signs or symptoms of liver inflammation or increasing conjugated bilirubin, alkaline phosphatase, or INR.

Coadministration with Drugs Metabolised by CYP3A

Also refer to CONTRAINDICATIONS, INTERACTIONS WITH OTHER MEDICINES, Table 13 and PHARMACOKINETICS-Implications for Drug Interactions.

Use with Fluticasone (glucocorticoids metabolised by CYP3A)

Use caution when administering VIEKIRA PAK with fluticasone or other glucocorticoids that are metabolised by CYP3A4. Concomitant use of inhaled glucocorticoids metabolised with CYP3A can increase systemic exposures of the glucocorticoids and cases of Cushing's syndrome and subsequent adrenal suppression have been reported with ritonavir-containing regimens. Concomitant use of VIEKIRA PAK and glucocorticoids, particularly long-term use, should only be initiated if the potential benefit of treatment outweighs the risk of systemic corticosteroid effects.

Use with Quetiapine

The use of VIEKIRA PAK with quetiapine is not recommended due to increases in quetiapine exposure. If coadministration is necessary, reduce the quetiapine dose to 1/6th of the current dose and monitor for quetiapine-associated adverse reactions. Refer to the quetiapine prescribing information for the recommendations on adverse reaction monitoring.

Use with colchicine

The interaction between VIEKIRA PAK and colchicine has not been evaluated. A reduction in colchicine dosage or an interruption of colchicine treatment is recommended in patients with normal renal or hepatic function if treatment with VIEKIRA PAK is required. In patients with renal or hepatic impairment, use of colchicine with VIEKIRA PAK is contraindicated (see CONTRAINDICATIONS).

Use with statins

Simvastatin and lovastatin are contraindicated (see CONTRAINDICATIONS).

Atorvastatin, Pitavastatin and fluvastatin

The interactions between atorvastatin, pitavastatin and fluvastatin and VIEKIRA PAK have not been investigated. Theoretically, VIEKIRA PAK is expected to increase the exposure to atorvastatin, pitavastatin and fluvastatin. A temporary suspension of atorvastatin, pitavastatin, fluvastatin is recommended for the duration of treatment with VIEKIRA PAK. If statin treatment is required during the treatment period, a switch to a reduced dose of pravastatin or rosuvastatin is possible (see Table 13).

Hepatic Impairment

No dose adjustment of VIEKIRA PAK is required in patients with mild hepatic impairment (Child-Pugh A). The safety and efficacy of VIEKIRA PAK have not been established in HCV-infected patients with moderate hepatic impairment (Child-Pugh B); however, no dose adjustment is expected to be required based on pharmacokinetic studies. VIEKIRA PAK is contraindicated in patients with severe hepatic impairment (Child-Pugh C) (see CONTRAINDICATIONS and PRECAUTIONS).

Treatment of Patients with Other HCV Genotypes

The safety and efficacy of VIEKIRA PAK has not been established in patients with HCV genotypes other than genotype 1.

Effects on Fertility

Paritaprevir/ritonavir

Paritaprevir/ritonavir had no effects on embryofetal viability or on fertility when evaluated in rats up to the highest dose of 300/30 mg/kg/day. Paritaprevir and ritonavir AUC exposures at this dosage were approximately 2 and 3-fold the exposure in humans at the recommended clinical dose.

Ombitasvir

Ombitasvir had no effects on embryofetal viability or on fertility when evaluated in mice up to the highest dose of 200 mg/kg/day. Ombitasvir AUC exposures at this dosage were approximately 23-fold (female) or 29-fold (male) the exposure in humans at the recommended clinical dose.

Dasabuvir

Dasabuvir had no effects on embryofetal viability or on fertility when evaluated in rats up to the highest dosage of 800 mg/kg/day. Dasabuvir AUC exposures at this dosage were approximately 16-fold the exposure in humans at the recommended clinical dose.

Use in Pregnancy

Pregnancy Category B3: VIEKIRA PAK

Since there are no adequate and well-controlled studies with VIEKIRA PAK in pregnant women, it should be used during pregnancy only if the benefits outweigh the risks.

No effects on embryofetal development have been noted in studies in animals with paritaprevir/ritonavir (in combination), ombitasvir and its major inactive human metabolites (M29, M36) or dasabuvir. For paritaprevir/ritonavir, the highest doses tested produced exposures equal to 98-fold (mouse) or 8-fold (rat) (for paritaprevir and 8-fold (mouse) or 3-fold (rat) for ritonavir) the exposures in humans at the recommended clinical doses. For ombitasvir, the highest dose tested produced exposures equal to 28-fold (mouse) or 4-fold (rabbit) the exposures in humans at the recommended clinical dose. The highest doses of the major, inactive human metabolites similarly tested produced exposures approximately 26 times higher in mice than in humans at the recommended clinical dose.

Developmental toxicity has been observed in embryofetal development studies with ritonavir alone. In rats, early resorptions, decreased fetal body weight and ossification delays and developmental variations occurred at a maternally toxic dosage of 75 mg/kg/day (5-fold the exposure in humans at the recommended clinical dose). A slight increase in the incidence of cryptorchidism was also noted in rats given 35 mg/kg/day (4-fold the exposure in humans at the recommended clinical dose). Developmental toxicity observed in rabbits (resorptions, decreased litter size and decreased fetal weights) also occurred at a maternally toxic dosage of 110 mg/kg/day. For dasabuvir, the highest dose tested produced exposures equal to 24-fold (rat) or 6-fold (rabbit) the exposures in humans at the recommended clinical dose. Developmental effects have not been identified in humans exposed to ritonavir during pregnancy nor has there been an association with cryptorchidism.

Use in Lactation

It is not known whether paritaprevir/ritonavir, ombitasvir, dasabuvir and their metabolites or ribavirin are excreted in human breast milk. Paritaprevir and its hydrolysis product M13, unchanged ombitasvir and dasabuvir were the predominant components observed in the milk of lactating rats, without effect on nursing pups.

Because of the potential for adverse reactions in nursing infants, breast feeding must be discontinued prior to initiation of treatment.

Paediatric Use

Safety and effectiveness of VIEKIRA PAK in children less than 18 years of age have not been established.

Use in the Elderly

No dose adjustment of VIEKIRA PAK is warranted in elderly patients. In Phase 3 clinical trials, 8.5% (174/2053) of subjects were age 65 or over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger subjects. The safety and effectiveness of VIEKIRA PAK has not been established in patients aged 70 years or over.

Genotoxicity

Paritaprevir

Paritaprevir was positive in an *in vitro* human chromosome aberration test. Paritaprevir was negative in a bacterial mutation assay, and in two *in vivo* genetic toxicology assays (rat bone marrow micronucleus and rat liver Comet tests).

Ombitasvir

Ombitasvir and its major inactive human metabolites (M29, M36) were not genotoxic in a battery of *in vitro* or *in vivo* assays, including bacterial mutagenicity, chromosome aberration using human peripheral blood lymphocytes and *in vivo* mouse micronucleus assays.

Dasabuvir

Dasabuvir was not genotoxic in a battery of *in vitro* or *in vivo* assays, including bacterial mutagenicity, chromosome aberration using human peripheral blood lymphocytes and *in vivo* rat micronucleus assays.

Ritonavir

Ritonavir showed no mutagenic potential in a series of assays for gene mutations (*S. typhimurium*, *E. coli* and mouse lymphoma cells) and chromosomal damage (mouse micronucleus assay *in-vivo* and human lymphocytes *in-vitro*).

Carcinogenicity

Paritaprevir/ritonavir was not carcinogenic in a 6-month transgenic mouse study up to the highest dosage tested (300/30 mg/kg/day), resulting in paritaprevir and ritonavir AUC exposures approximately 38 and 5-fold higher, respectively than those in humans at the recommended dose of 150/50 mg. Similarly, paritaprevir/ritonavir was not carcinogenic in a 2-year rat study up to the highest dosage tested (300/30 mg/kg/day), resulting in paritaprevir/ritonavir AUC exposures approximately 8/5-fold higher than those in humans at 150/50 mg.

Two-year carcinogenicity studies have been conducted in rodents with ritonavir alone at dietary levels of 50, 100 and 200 mg/kg/day in mice, and 7, 15 and 30 mg/kg/day in rats. In male mice there was a dose-dependent increase in the incidence of hepatocellular adenomas, and adenomas and carcinomas combined, both reaching statistical significance only at the high-dose. In female mice there were small, statistically significant increases in these tumour incidences only at the high-dose. In rats, there were no tumorigenic effects.

Ombitasvir was not carcinogenic in a 6-month transgenic mouse study up to the highest dosage tested (150 mg/kg/day), resulting in ombitasvir AUC exposures approximately 26-fold higher than those in humans at the recommended clinical dose of 25 mg. The carcinogenicity study of ombitasvir in rats is ongoing.

Dasabuvir was not carcinogenic in a 6-month transgenic mouse study up to the highest dosage tested (2000 mg per kg per day), resulting in dasabuvir AUC exposures approximately 19-fold higher than those in humans at the recommended dose of 500 mg (250 mg twice daily).

The carcinogenicity study of dasabuvir in rats is ongoing.

Effect on Laboratory Tests

Changes in selected laboratory parameters are described in Table 12. A side-by-side tabulation is provided to simplify presentation; direct comparisons should not be made across trials that differ in study design.

Table 12: Selected Treatment Emergent Laboratory Abnormalities

Laboratory Parameters	SAPPHIRE I and II		PEARL II, III and IV		TURQUOISE II (subjects with cirrhosis)
	VIEKIRA PAK + RBV 12 Weeks N = 770 n (%)	Placebo 12 Weeks N = 255 n (%)	VIEKIRA PAK + RBV 12 Weeks N = 401 n (%)	VIEKIRA PAK 12 Weeks N = 509 n (%)	VIEKIRA PAK + RBV 12 or 24 Weeks N = 380 n (%)
ALT					
> 5-20 × ULN* (Grade 3)	6/765 (0.8%)	10/254 (3.9%)	3/401 (0.7%)	1/509 (0.2%)	4/380 (1.1%)
> 20 × ULN (Grade 4)	3/765 (0.4%)	0	0	0	2/380 (0.5%)
Haemoglobin					
< 10-8 g/dL (Grade 2)	41/765 (5.4%)	0	23/401 (5.7%)	0	30/380 (7.9%)
< 8-6.5 g/dL (Grade 3)	1/765 (0.1%)	0	2/401 (0.5%)	0	3/380 (0.8%)
< 6.5 g/dL (Grade 4)	0	0	0	0	1/380 (0.3%)
Total Bilirubin					
> 3-10 × ULN (Grade 3)	19/765 (2.5%)	0	23/401 (5.7%)	2/509 (0.4%)	37/380 (9.7%)
> 10 × ULN (Grade 4)	1/765 (0.1%)	0	0	0	0
*ULN: Upper Limit of Normal according to testing laboratory.					

Serum ALT elevations

During clinical trials with VIEKIRA PAK and VIEKIRA PAK-RBV, less than 1% of subjects who were not on systemic ethinyl estradiol-containing medications experienced transient serum ALT levels greater than 5 times the upper limit of normal (ULN) after starting treatment. These elevations were asymptomatic, generally occurred during the first 4 weeks of treatment and resolved with ongoing therapy. Elevations in ALT were generally not associated with bilirubin elevations. Cirrhosis was not a risk factor for elevated ALT (see PRECAUTIONS).

Serum Bilirubin Elevations

Transient elevations in bilirubin (predominantly indirect) were observed in subjects receiving VIEKIRA PAK, related to the inhibition of the bilirubin transporters OATP1B1/1B3 by paritaprevir and ribavirin-induced haemolysis. Bilirubin elevations occurred after initiation of treatment, peaked by study Week 1, and generally resolved with ongoing therapy. Bilirubin elevations were not associated with aminotransferase elevations. The frequency of indirect bilirubin elevations was lower among subjects who did not receive ribavirin.

INTERACTIONS WITH OTHER MEDICINES

Also refer to the CONTRAINDICATIONS, INTERACTIONS WITH OTHER MEDICINES SECTION, Table 13 and PHARMACOKINETICS-Implications for Drug Interactions sections.

Recommendations for co-administration of VIEKIRA PAK for a number of medicinal products are provided in Table 13.

If a patient is already taking medicinal product(s) or initiating a medicinal product while receiving VIEKIRA PAK for which potential for drug interaction is expected, dose adjustment of the concomitant medicinal product(s) or appropriate clinical monitoring should be considered (Table 13).

A change of 0.5- to 2.0-fold in the exposures (C_{max} and AUC) of paritaprevir, ombitasvir and dasabuvir is not considered clinically relevant and does not require dose adjustment for VIEKIRA PAK.

If dose adjustments of concomitant medicinal products are made due to treatment with VIEKIRA PAK, doses should be re-adjusted after administration of VIEKIRA PAK is completed.

Table 13 provides the Least Squares Means Ratio (90% Confidence Interval) effect on concentration of VIEKIRA PAK and concomitant medicinal products. Dose adjustment is not required for VIEKIRA PAK when administered with the concomitant medications listed in Table 13 unless otherwise noted.

Table 13: Interactions between VIEKIRA PAK and other medicinal products

Drug Class Drug Name	Effect	Clinical Comment
Aminosalicylate		
Sulfasalazine	↑ sulfasalazine*	Mechanism: BCRP inhibition by paritaprevir, ritonavir and dasabuvir. Caution should be used when sulfasalazine is co-administered with VIEKIRA PAK.
Angiotensin receptor blocker		

Attachment 1: Product information AusPAR VIEKIRA PAK - Paritaprevir / Ritonavir / Ombitasvir and Dasabuvir (as Sodium Salt) - AbbVie Pty Ltd - PM-2014-01436-1-2 FINAL 24 March 2017. This Product Information was approved at the time this AusPAR was published.

Valsartan	↑ valsartan*			Mechanism: OATP1B inhibition by paritaprevir. Clinical monitoring and dose reduction is recommended when VIEKIRA PAK is coadministered with valsartan.	
Antiarrhythmics					
Digoxin		C_{max}	AUC	C_{min}	Mechanism: P-gp inhibition by paritaprevir, ritonavir and dasabuvir. While no dose adjustment is necessary for digoxin, appropriate monitoring of serum digoxin levels is recommended.
	↔ digoxin	1.15 (1.04-1.27)	1.16 (1.09-1.23)	1.01 (0.97-1.05)	
	↔ ombitasvir	1.03 (0.97-1.10)	1.00 (0.98-1.03)	0.99 (0.96-1.02)	
	↔ paritaprevir	0.92 (0.80-1.06)	0.94 (0.81-1.08)	0.92 (0.82-1.02)	
	↔ dasabuvir	0.99 (0.92-1.07)	0.97 (0.91-1.02)	0.99 (0.92-1.07)	
Antibiotics					
Erythromycin	↑ erythromycin*			Mechanism: CYP3A4/P-gp inhibition by paritaprevir, ritonavir and dasabuvir. Caution is advised when erythromycin is administered with Viekira Pak.	
Anticancer agents					
Imatinib	↑ imatinib*			Mechanism: BCRP inhibition by paritaprevir, ritonavir and dasabuvir. Clinical monitoring and lower doses of imatinib are recommended.	
Anticoagulants					
Warfarin 5 mg single dose		C_{max}	AUC	C_{min}	While no dose adjustment is necessary for warfarin, appropriate monitoring of international normalised ratio (INR) is recommended.
	↔ R-warfarin	1.05 (0.95-1.17)	0.88 (0.81-0.95)	0.94 (0.84-1.05)	
	↔ S-warfarin	0.96 (0.85-1.08)	0.88 (0.81-0.96)	0.95 (0.88-1.02)	
	↔ ombitasvir	0.94 (0.89-1.00)	0.96 (0.93-1.00)	0.98 (0.95-1.02)	
	↔ paritaprevir	0.98 (0.82-1.18)	1.07 (0.89-1.27)	0.96 (0.85-1.09)	
↔ dasabuvir	0.97 (0.89-1.06)	0.98 (0.91-1.06)	1.03 (0.94-1.13)		
Dabigatran etexilate	↑ dabigatran*			Mechanism: Intestinal P-gp inhibition by paritaprevir, dasabuvir and ritonavir. VIEKIRA PAK may increase the plasma concentrations of dabigatran etexilate. Use with caution.	
S-mephenytoin	↓ S-mephenytoin*			Mechanism: CYP2C19 induction by ritonavir. Clinical monitoring and dose adjustment maybe needed for s-mephenytoin.	
Antidepressants					
Escitalopram 10 mg single dose		C_{max}	AUC	C_{min}	No dose adjustment is necessary for escitalopram.
	↔ es-citalopram	1.00 (0.96-1.05)	0.87 (0.80-0.95)	NA	
	↑ S-Desmethyl citalopram	1.15 (1.10-1.21)	1.36 (1.03-1.80)	NA	

Attachment 1: Product information AusPAR VIEKIRA PAK - Paritaprevir / Ritonavir / Ombitasvir and Dasabuvir (as Sodium Salt) - AbbVie Pty Ltd - PM-2014-01436-1-2 FINAL 24 March 2017. This Product Information was approved at the time this AusPAR was published.

	↔ ombitasvir	1.09 (1.01-1.18)	1.02 (1.00-1.05)	0.97 (0.92-1.02)	
	↔ paritaprevir	1.12 (0.88-1.43)	0.98 (0.85-1.14)	0.71 (0.56-0.89)	
	↔ dasabuvir	1.10 (0.95-1.27)	1.01 (0.93-1.10)	0.89 (0.79-1.00)	
Duloxetine 60 mg single dose		C_{max}	AUC	C_{min}	No dose adjustment is necessary for duloxetine.
	↓ duloxetine	0.79 (0.67-0.94)	0.75 (0.67-0.83)	NA	
	↔ ombitasvir	0.98 (0.88-1.08)	1.00 (0.95-1.06)	1.01 (0.96-1.06)	
	↓ paritaprevir	0.79 (0.53-1.16)	0.83 (0.62-1.10)	0.77 (0.65-0.91)	
	↔ dasabuvir	0.94 (0.81-1.09)	0.92 (0.81-1.04)	0.88 (0.76-1.01)	

Attachment 1: Product information AusPAR VIEKIRA PAK - Paritaprevir / Ritonavir / Ombitasvir and Dasabuvir (as Sodium Salt) - AbbVie Pty Ltd - PM-2014-01436-1-2 FINAL 24 March 2017. This Product Information was approved at the time this AusPAR was published.

Trazodone	↑ Trazodone*				Mechanism: CYP3A4 inhibition by ritonavir. Trazodone should be used with caution and a lower dose of trazodone may be considered.
Antifungals					
Ketoconazole 400 mg once daily.		C _{max}	AUC	C _{min}	When VIEKIRA PAK is co-administered with ketoconazole, the maximum daily dose of ketoconazole should be limited to 200 mg per day.
	↑ ketoconazole	1.15 (1.09-1.21)	2.17 (2.05-2.29)	NA	
	↔ ombitasvir	0.98 (0.90-1.06)	1.17 (1.11-1.24)	NA	
	↑ paritaprevir	1.37 (1.11-1.69)	1.98 (1.63-2.42)	NA	
	↑ dasabuvir	1.16 (1.03-1.32)	1.42 (1.26-1.59)	NA	
Voriconazole	↓ voriconazole*				Mechanism: CYP2C19 induction by ritonavir. Co-administration of VIEKIRA PAK with voriconazole is not recommended unless an assessment of the benefit-to-risk ratio justifies the use of voriconazole.
Antihistamines (for systemic use)					
Fexofenadine	↑ fexofenadine*				Mechanism: OATP1B1 inhibition by paritaprevir. Caution should be used when VIEKIRA PAK is coadministered with fexofenadine.
Antihyperlipidemic agent					
Gemfibrozil		C _{max}	AUC	C _{min}	10-fold increase in dasabuvir exposure. Increased risk of QT-prolongation (see CONTRAINDICATIONS).
	↑ paritaprevir	1.21 (0.94-1.57)	1.38 (1.18-1.61)	NA	
	↑ dasabuvir	2.01 (1.71-2.38)	11.25 (9.05-13.99)	NA	
Calcium Channel Blockers					
Amlodipine 5 mg single dose		C _{max}	AUC	C _{min}	Mechanism: CYP3A4 inhibition by ritonavir. Decrease amlodipine dose by 50% and monitor patients for clinical effects.
	↑ amlodipine	1.26 (1.11-1.44)	2.57 (2.31-2.86)	NA	
	↔ ombitasvir	1.00 (0.95-1.06)	1.00 (0.97-1.04)	1.00 (0.97-1.04)	
	↓ paritaprevir	0.77 (0.64-0.94)	0.78 (0.68-0.88)	0.88 (0.80-0.95)	
	↔ dasabuvir	1.05 (0.97-1.14)	1.01 (0.96-1.06)	0.95 (0.89-1.01)	
Diltiazem Verapamil	↑ diltiazem*, verapamil* ↑ paritaprevir ↑/↔ dasabuvir				Mechanism: CYP3A4/P-gp inhibition Caution is advised due to the expected increase in paritaprevir exposures. Dose decrease and clinical monitoring of calcium channel blockers is recommended when co-administered with VIEKIRA PAK.
Nifedipine	↑ nifedipine*				Mechanism: CYP3A4 inhibition Dose decrease and clinical monitoring of calcium channel blockers is recommended when co-administered with VIEKIRA PAK.

Attachment 1: Product information AusPAR VIEKIRA PAK - Paritaprevir / Ritonavir / Ombitasvir and Dasabuvir (as Sodium Salt) - AbbVie Pty Ltd - PM-2014-01436-1-2 FINAL 24 March 2017. This Product Information was approved at the time this AusPAR was published.

Diuretics					
Furosemide 20 mg single dose		C_{max}	AUC	C_{min}	Mechanism: Possibly due to UGT1A1 inhibition by paritaprevir, ombitasvir and dasabuvir. Patients should be monitored for clinical effects; a decrease in furosemide dose of up to 50% may be required.
	↑ furosemide	1.42 (1.17-1.72)	1.08 (1.00-1.17)	NA	
	↔ ombitasvir	1.14 (1.03-1.26)	1.07 (1.01-1.12)	1.12 (1.08-1.16)	
	↔ paritaprevir	0.93 (0.63-1.36)	0.92 (0.70-1.21)	1.26 (1.16-1.38)	
	↔ dasabuvir	1.12 (0.96-1.31)	1.09 (0.96-1.23)	1.06 (0.98-1.14)	
HIV Antivirals: Protease Inhibitors					
Atazanavir 300 mg once daily (given at the same time)		C_{max}	AUC	C_{min}	Mechanism: Increase in paritaprevir exposures may be due to inhibition of OATP1B1/B3 and CYP3A by atazanavir. In combination with ritonavir, the increase in paritaprevir exposures may be due to inhibition of OATP1B1/B3 and CYP3A by atazanavir and CYP3A by the additional dose of ritonavir.
	↔ atazanavir	0.91 (0.84-0.99)	1.01 (0.93-1.10)	0.90 (0.81-1.01)	
	↓ ombitasvir	0.77 (0.70-0.85)	0.83 (0.74-0.94)	0.89 (0.78-1.02)	
	↑ paritaprevir	1.46 (1.06-1.99)	1.94 (1.34-2.81)	3.26 (2.06-5.16)	
	↔ dasabuvir	0.83 (0.71-0.96)	0.82 (0.71-0.94)	0.79 (0.66-0.94)	
Atazanavir/ ritonavir 300/100 mg once daily (administered 12 hours apart)		C_{max}	AUC	C_{min}	The recommended dose of atazanavir is 300 mg, without ritonavir, in combination with VIEKIRA PAK. Atazanavir must be administered at the same time as VIEKIRA PAK. Ritonavir dose in VIEKIRA PAK will provide atazanavir pharmacokinetic enhancement. Treatment with atazanavir/ritonavir + VIEKIRA PAK is not recommended.
	↔ atazanavir	1.02 (0.92-1.13)	1.19 (1.11-1.28)	1.68 (1.44-1.95)	
	↔ ombitasvir	0.83 (0.72-0.96)	0.90 (0.78-1.02)	1.00 (0.89-1.13)	
	↑ paritaprevir	2.19 (1.61-2.98)	3.16 (2.40-4.17)	11.95 (8.94-15.98)	
	↔ dasabuvir	0.81 (0.73-0.91)	0.81 (0.71-0.92)	0.80 (0.65-0.98)	
Darunavir 800 mg once daily (given at the same time)		C_{max}	AUC	C_{min}	The recommended dose of darunavir is 800 mg once daily, without ritonavir, when administered at the same time as VIEKIRA PAK (ritonavir dose in Viekira Pak will provide darunavir pharmacokinetic enhancement). This regimen can be used in the absence of extensive PI resistance (i.e. lack of darunavir associated RAMs). Darunavir combined with VIEKIRA PAK is not recommended in patients with extensive PI resistance.
	↓ darunavir	0.92 (0.87-0.98)	0.76 (0.71-0.82)	0.52 (0.47-0.58)	
	↔ ombitasvir	0.86 (0.77-0.95)	0.86 (0.79-0.94)	0.87 (0.82-0.92)	
	↑ paritaprevir	1.54 (1.14-2.09)	1.29 (1.04-1.61)	1.30 (1.09-1.54)	
	↔ dasabuvir	1.10 (0.88-1.37)	0.94 (0.78-1.14)	0.90 (0.76-1.06)	
Darunavir/ ritonavir 600/100 mg twice daily		C_{max}	AUC	C_{min}	
	↔ darunavir	0.87 (0.79-0.96)	0.80 (0.74-0.86)	0.57 (0.48-0.67)	
	↓ ombitasvir	0.76 (0.65-0.88)	0.73 (0.66-0.80)	0.73 (0.64-0.83)	
	↓ paritaprevir	0.70 (0.43-1.12)	0.59 (0.44-0.79)	0.83 (0.69-1.01)	
	↓ dasabuvir	0.84 (0.67-1.05)	0.73 (0.62-0.86)	0.54 (0.49-0.61)	
Darunavir/ ritonavir 800/100 mg once daily (administered 12 hours apart)		C_{max}	AUC	C_{min}	
	↑ darunavir	0.79 (0.70-0.90)	1.34 (1.25-1.43)	0.54 (0.48-0.62)	
	↔ ombitasvir	0.87 (0.82-0.93)	0.87 (0.81-0.93)	0.87 (0.80-0.95)	
	↓ paritaprevir	0.70 (0.50-0.99)	0.81 (0.60-1.09)	1.59 (1.23-2.05)	
	↓ dasabuvir	0.75 (0.64-0.88)	0.72 (0.64-0.82)	0.65 (0.58-0.72)	
Lopinavir / ritonavir 400/100 mg twice daily ²		C_{max}	AUC	C_{min}	Mechanism: Increase in paritaprevir exposures may be due to inhibition of CYP3A/efflux transporters by lopinavir and higher dose of ritonavir.
	↔ lopinavir	0.87 (0.76-0.99)	0.94 (0.81-1.10)	1.15 (0.93-1.42)	
	↔ ombitasvir	1.14 (1.01-1.28)	1.17 (1.07-1.28)	1.24 (1.14-1.34)	

Attachment 1: Product information AusPAR VIEKIRA PAK - Paritaprevir / Ritonavir / Ombitasvir and Dasabuvir (as Sodium Salt) - AbbVie Pty Ltd - PM-2014-01436-1-2 FINAL 24 March 2017. This Product Information was approved at the time this AusPAR was published.

	↑ paritaprevir	2.04 (1.30-3.20)	2.17 (1.63-2.89)	2.36 (1.00-5.55)	Lopinavir/ritonavir 400/100 mg twice daily and 800/200 mg once daily (evening administration) increases paritaprevir concentrations. Lopinavir/ritonavir use is not recommended with VIEKIRA PAK.
	↔ dasabuvir	0.99 (0.75-1.31)	0.93 (0.75-1.15)	0.68 (0.57-0.80)	
HIV Antivirals: Non-nucleoside reverse transcriptase inhibitors					
Rilpivirine 25 mg once daily administered in the morning, with food ³		C_{max}	AUC	C_{min}	Mechanism: CYP3A4 inhibition by ritonavir. Co-administration of VIEKIRA PAK with rilpivirine once daily is not recommended due to potential for QT interval prolongation with higher concentrations of rilpivirine.
	↑ rilpivirine	2.55 (2.08-3.12)	3.25 (2.80-3.77)	3.62 (3.12-4.21)	
	↔ ombitasvir	1.11 (1.02-1.20)	1.09 (1.04-1.14)	1.05 (1.01-1.08)	
	↑ paritaprevir	1.30 (0.94-1.81)	1.23 (0.93-1.64)	0.95 (0.84-1.07)	
	↔ dasabuvir	1.18 (1.02-1.37)	1.17 (0.99-1.38)	1.10 (0.89-1.37)	
Efavirenz/emtricitabine/tenofovir disoproxil fumarate 600/300/200 mg once daily	Co-administration of efavirenz (enzyme inducer) based regimens with paritaprevir/ritonavir + dasabuvir resulted in ALT elevations and therefore, early discontinuation of the study.			Co-administration of efavirenz based regimens with paritaprevir, ritonavir plus dasabuvir was poorly tolerated and resulted in liver enzyme elevations. (see CONTRAINDICATIONS).	
HIV Antivirals: Integrase strand transfer inhibitor					
Raltegravir 400 mg twice daily		C_{max}	AUC	C_{min}	No dose adjustment is necessary for raltegravir.
	↑ raltegravir	2.33 (1.66-3.27)	2.34 (1.70-3.24)	2.00 (1.17-3.42)	
HIV ANTIVIRALS: NUCLEOSIDE INHIBITORS					
Emtricitabine/tenofovir 200 mg once daily/300 mg once daily	↔ em-tricitabine	1.05 (1.00-1.12)	1.07 (1.00-1.14)	1.09 (1.01-1.17)	No dose adjustment is necessary for emtricitabine/tenofovir.
	↔ tenofovir	1.07 (0.93-1.24)	1.13 (1.07-1.20)	1.24 (1.13-1.36)	
	↔ ombitasvir	0.89 (0.81-0.97)	0.99 (0.93-1.05)	0.97 (0.90-1.04)	
	↓ paritaprevir	0.68 (0.42-1.11)	0.84 (0.59-1.17)	1.06 (0.83-1.35)	
	↔ dasabuvir	0.85 (0.74-0.98)	0.85 (0.75-0.96)	0.85 (0.73-0.98)	
HMG-CoA Reductase Inhibitors					
Rosuvastatin 5 mg once daily		C_{max}	AUC	C_{min}	Mechanism: OATP1B inhibition by paritaprevir and BCRP inhibition by paritaprevir, ritonavir or dasabuvir The maximum daily dose of rosuvastatin should be 5 mg
	↑ rosuvastatin	7.13 (5.11-9.96)	2.59 (2.09-3.21)	0.59 (0.51-0.69)	
	↔ ombitasvir	0.92 (0.82-1.04)	0.89 (0.83-0.95)	0.88 (0.83-0.94)	
	↑ paritaprevir	1.59 (1.13-2.23)	1.52 (1.23-1.90)	1.43 (1.22-1.68)	
	↔ dasabuvir	1.07 (0.92-1.24)	1.08 (0.92-1.26)	1.15 (1.05-1.25)	
Pravastatin 10 mg once daily		C_{max}	AUC	C_{min}	Mechanism: OATP1B/CYP3A4 inhibition by paritaprevir. Reduce pravastatin dose by 50%.
	↑ pravastatin	1.37 (1.11-1.69)	1.82 (1.60-2.08)	NA	
	↔ ombitasvir	0.95 (0.89-1.02)	0.89 (0.83-0.95)	0.94 (0.89-0.99)	
	↔ dasabuvir	1.00 (0.87-1.14)	0.96 (0.85-1.09)	1.03 (0.91-1.15)	
	↔ paritaprevir	0.96 (0.69-1.32)	1.13 (0.92-1.38)	1.39 (1.21-1.59)	

Attachment 1: Product information AusPAR VIEKIRA PAK - Paritaprevir / Ritonavir / Ombitasvir and Dasabuvir (as Sodium Salt) - AbbVie Pty Ltd - PM-2014-01436-1-2 FINAL 24 March 2017. This Product Information was approved at the time this AusPAR was published.

Immunosuppressants					
Ciclosporin 30 mg once daily single dose ⁴		C_{max}	AUC	C_{min}	When starting co-administration with VIEKIRA PAK, give one fifth of the total daily dose of ciclosporin once daily with VIEKIRA PAK. Monitor ciclosporin levels and adjust dose and/or dosing frequency as needed.
	↑ cilosporin	1.01 (0.85-1.20)	5.82 (4.73-7.14)	15.8 (13.8-18.09)	
	↔ ombitasvir	0.99 (0.92-1.07)	1.08 (1.05-1.11)	1.15 (1.08-1.23)	
	↑ paritaprevir	1.44 (1.16-1.78)	1.72 (1.49-1.99)	1.85 (1.58-2.18)	
	↓ dasabuvir	0.66 (0.58-0.75)	0.70 (0.65-0.76)	0.76 (0.71-0.82)	
Tacrolimus 2 mg single dose		C_{max}	AUC	C_{min}	Mechanism: Effect on tacrolimus is due to CYP3A4 inhibition by ritonavir. When starting co-administration with VIEKIRA PAK, administer 0.5 mg tacrolimus once every week. Monitor tacrolimus levels and adjust dose and/or dosing frequency as needed.
	↑ tacrolimus	3.99 (3.21-4.97)	57.1 (45.5-71.7)	16.6 (13.0-21.2)	
	↔ ombitasvir	0.93 (0.88-0.99)	0.94 (0.89-0.98)	0.94 (0.91-0.96)	
	↓ paritaprevir	0.57 (0.42-0.78)	0.66 (0.54-0.81)	0.73 (0.66-0.80)	
	↔ dasabuvir	0.85 (0.73-0.98)	0.90 (0.80-1.02)	1.01 (0.91-1.11)	
Insulin Secretagogues					
Repaglinide	↑ repaglinide*				Mechanism: OATP1B1 inhibition by paritaprevir. Caution should be used and dose decrease maybe needed for repaglinide
Iron Chelators					
Deferasirox	↑ dasabuvir*				Mechanism: CYP2C8 inhibition by deferasirox. Deferasirox may increase dasabuvir exposures and should be used with caution.
Medicinal Products used in Multiple Sclerosis					
Teriflunomide	↑ dasabuvir*				CYP2C8 inhibition by teriflunomide. Teriflunomide may increase dasabuvir exposures and should be used with caution.
Opioids					
Methadone 20-120 mg once daily ⁵		C_{max}	AUC	C_{min}	No dose adjustment is necessary for methadone.
	↔ R- Methadone	1.04 (0.98-1.11)	1.05 (0.98-1.11)	0.94 (0.87-1.01)	
	↔ S- Methadone	0.99 (0.91-1.08)	0.99 (0.89-1.09)	0.86 (0.76-0.96)	
	↔ paritaprevir/ombitasvir/dasabuvir (based on the cross-study comparison)				
Buprenorphine/ naloxone 4-24 mg/1-6 mg once daily ⁵		C_{max}	AUC	C_{min}	No dose adjustment is necessary for buprenorphine/naloxone.
	↑ bu- prenorphine	2.18 (1.78-2.68)	2.07 (1.78-2.40)	3.12 (2.29-4.27)	
	↑ norbu- prenorphine	2.07 (1.42-3.01)	1.84 (1.30-2.60)	2.10 (1.49-2.97)	
	↑ naloxone	1.18 (0.81-1.73)	1.28 (0.92-1.79)	NA	
	↔ ombitasvir/paritaprevir/dasabuvir (based on the cross-study comparison)				
Proton Pump Inhibitors					
Omeprazole 40 mg once daily		C_{max}	AUC	C_{min}	Mechanism: CYP2C19 induction by ritonavir. If clinically indicated higher doses of omeprazole should be used.
	↓ omeprazole	0.62 (0.48-0.80)	0.62 (0.51-0.75)	NA	
	↔ ombitasvir	1.02 (0.95-1.09)	1.05 (0.98-1.12)	1.04 (0.98-1.11)	
	↔ paritaprevir	1.19 (1.04-1.36)	1.18 (1.03-1.37)	0.92 (0.76-1.12)	
	↔ dasabuvir	1.13 (1.03-1.25)	1.08 (0.98-1.20)	1.05 (0.93-1.19)	

Attachment 1: Product information AusPAR VIEKIRA PAK - Paritaprevir / Ritonavir / Ombitasvir and Dasabuvir (as Sodium Salt) - AbbVie Pty Ltd - PM-2014-01436-1-2 FINAL 24 March 2017. This Product Information was approved at the time this AusPAR was published.

Esomeprazole, Lansoprazole	↓ esomeprazole*, lansoprazole*	Mechanism: CYP2C19 induction by ritonavir . If clinically indicated, higher doses of esomeprazole/lansoprazole may be needed.			
Sedatives/hypnotics					
Triazolam, orally administered midazolam	Large ↑ triazolam, orally administered midazolam	Triazolam and orally administered midazolam are extensively metabolised by CYP3A4. Coadministration of triazolam or orally administered midazolam with VIEKIRA PAK may cause large increases in the concentration of these benzodiazepines. The potential exists for serious and/or life threatening events such as prolonged or increased sedation or respiratory depression. (see CONTRAINDICATIONS).			
Zolpidem 5 mg single dose		C _{max}	AUC	C _{min}	No dose adjustment is necessary for zolpidem.
	↔ zolpidem	0.94 (0.76-1.16)	0.95 (0.74-1.23)	NA	
	↔ ombitasvir	1.07 (1.00-1.15)	1.03 (1.00-1.07)	1.04 (1.00-1.08)	
	↓ paritaprevir	0.63 (0.46-0.86)	0.68 (0.55-0.85)	1.23 (1.10-1.38)	
	↔ dasabuvir	0.93 (0.84-1.03)	0.95 (0.84-1.08)	0.92 (0.83-1.01)	
Alprazolam 0.5 mg single dose		C _{max}	AUC	C _{min}	Clinical monitoring of patients is recommended. A decrease in alprazolam dose can be considered based on clinical response.
	↑ alprazolam	1.09 (1.03-1.15)	1.34 (1.15-1.55)	NA	
	↔ ombitasvir	0.98 (0.93-1.04)	1.00 (0.96-1.04)	0.98 (0.93-1.04)	
	↔ paritaprevir	0.91 (0.64-1.31)	0.96 (0.73-1.27)	1.12 (1.02-1.23)	
	↔ dasabuvir	0.93 (0.83-1.04)	0.98 (0.87-1.11)	1.00 (0.87-1.15)	
Thyroid Hormones					
Levothyroxine	↑ levothyroxine*	Mechanism: UGT1A1 inhibition by paritaprevir, ombitasvir and dasabuvir. Clinical monitoring and dose adjustment may be required for levothyroxine.			
<p>* Not studied; expected effect.</p> <ol style="list-style-type: none"> 1. Drug interaction study carried out with paritaprevir/ritonavir + dasabuvir combination. 2. Lopinavir/ritonavir 800/200 mg once daily (administered in the evening) was also administered with VIEKIRA PAK. The effect on C_{max} and AUC of DAAs and lopinavir was similar to that observed when lopinavir/ritonavir 400/100 mg twice daily was administered with VIEKIRA PAK. 3. Rilpivirine was also administered in the evening with food and at night 4 hours after dinner with VIEKIRA PAK in other two arms in the study. The effect on rilpivirine exposures was similar to that observed when rilpivirine was administered in the morning with food with VIEKIRA PAK (shown in the table above). 4. Cyclosporin 100 mg dosed alone and 30 mg administered with VIEKIRA PAK. Dose normalized cyclosporine ratios are shown for interaction with VIEKIRA PAK. 5. Dose normalised parameters reported for methadone, buprenorphine and naloxone. <p>Note: Doses used for VIEKIRA PAK were: ombitasvir 25 mg, paritaprevir 150 mg, ritonavir 100 mg, once daily and dasabuvir 400 mg twice daily or 250 mg twice daily. The dasabuvir exposures obtained with the 400 mg formulation and the 250 mg tablet are similar. VIEKIRA PAK was administered as multiple doses in all the drug interaction studies except the drug interaction studies with carbamazepine, gemfibrozil and ketoconazole.</p>					

ADVERSE EFFECTS

Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in clinical trials of VIEKIRA PAK cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety summary is based on pooled data from phase 2 and 3 clinical trials in more than 2,600 subjects who received VIEKIRA PAK with or without ribavirin.

VIEKIRA PAK with Ribavirin in Subjects with Genotype 1 Hepatitis C Infection (including subjects with cirrhosis)

In subjects receiving VIEKIRA PAK, the most commonly reported adverse reactions (greater than 20% of subjects) were fatigue and nausea. The proportion of subjects who permanently discontinued treatment due to adverse events was 1.2% (25/2,044). 1.3% (27/2,044) of subjects interrupted treatment due to adverse events. 7.7% (158/2,044) of subjects had ribavirin dose reductions due to adverse events.

The safety profile of VIEKIRA PAK with ribavirin in subjects with cirrhosis was similar to that of subjects without cirrhosis.

VIEKIRA PAK without Ribavirin in Subjects with Genotype 1 Hepatitis C Infection

In subjects receiving VIEKIRA PAK without ribavirin, pruritus was the only identified adverse reaction (of reactions that occurred in $\geq 5\%$ of subjects) when a comparison of subjects who received VIEKIRA PAK without RBV was made to studies which included both VIEKIRA PAK with ribavirin and placebo.

The proportion of subjects who permanently discontinued treatment due to adverse events was 0.3% (2/588). 0.5% (3/588) subjects had treatment interruptions due to adverse events.

Table 13 lists adverse drug reactions from two randomised placebo-controlled trials (SAPPHIRE I and SAPPHIRE II) that occurred with at least 5% higher frequency among subjects receiving VIEKIRA PAK in combination with ribavirin compared to subjects receiving placebo, regardless of relationship to VIEKIRA PAK. In addition, Table 13 includes rates of these adverse events from three trials in which subjects received VIEKIRA PAK with or without ribavirin (PEARL II, PEARL III, and PEARL IV), and rates of these adverse events from the trial in subjects with cirrhosis who received VIEKIRA PAK or VIEKIRA PAK-RBV for 12 or 24 weeks (TURQUOISE II). A side-by-side tabulation is provided to simplify presentation; direct comparisons should not be made across trials that differ in design.

Table 14: Side-by-Side Tabulation of Adverse Event Rates in Phase 3 Trials Based on Adverse Reactions* (All Grades)

Adverse Reaction	SAPPHIRE I and II		PEARL II, III and IV		TURQUOISE II (subjects with cirrhosis)
	VIEKIRA PAK + RBV 12 Weeks N = 770 n (%)	Placebo 12 Weeks N = 255 n (%)	VIEKIRA PAK + RBV 12 Weeks N = 401 n (%)	VIEKIRA PAK 12 Weeks N = 509 n (%)	VIEKIRA PAK + RBV 12 or 24 Weeks N = 380 n (%)
Fatigue	263 (34.2)	67 (26.3)	120 (29.9)	135 (26.5)	148 (38.9)
Nausea	172 (22.3)	38 (14.9)	63 (15.7)	43 (8.4)	72 (18.9)
Pruritus	121 (15.7)	11 (4.3)	48 (12.0)	31 (6.1)**	71 (18.7)
Insomnia	108 (14.0)	19 (7.5)	49 (12.2)	26 (5.1)	63 (16.6)
Asthenia	104 (13.5)	17 (6.7)	36 (9.0)	20 (3.9)	51 (13.4)
Anaemia	41 (5.3)	0	30 (7.5)	1 (0.2)	34 (8.9)

*Adverse drug reactions for Viekira Pak with ribavirin listed are those with a 5% higher frequency among subjects receiving VIEKIRA PAK in combination with ribavirin compared to subjects receiving placebo in SAPPHIRE I and II.

** Adverse drug reaction for VIEKIRA PAK defined as the subset of ADRs for the VIEKIRA PAK- RBV for which the risk difference (VIEKIRA PAK-RBV minus VIEKIRA PAK) in PEARL II, III, and IV was at least 5.0 % lower than the risk difference (VIEKIRA PAK minus placebo) in SAPPHIRE I and II.

The majority of adverse events in the Phase 3 clinical trials were of grade 1 severity. The safety profile of VIEKIRA PAK with ribavirin was consistent with the known safety profile of ribavirin.

In addition to the adverse reaction listed in Table 14, treatment-emergent adverse events that occurred with at least 2% frequency and less than 5% higher frequency among subjects receiving VIEKIRA PAK-RBV compared to subjects receiving placebo (SAPPHIRE I and II), are listed below by system organ class.

<i>Gastrointestinal Disorders:</i>	Diarrhoea and vomiting
<i>Investigations:</i>	Haemoglobin decreased
<i>Metabolism and Nutrition Disorders:</i>	Decreased appetite
<i>Nervous System Disorders:</i>	Dizziness and headache
<i>Psychiatric Disorders:</i>	Sleep disorder
<i>Respiratory, Thoracic and Mediastinal Disorders:</i>	Cough and dyspnoea
<i>Skin and Subcutaneous Tissue Disorders:</i>	Dry skin, and rash

DOSAGE AND ADMINISTRATION

VIEKIRA PAK is paritaprevir/ritonavir/ombitasvir fixed dose combination tablets copackaged with dasabuvir tablets.

Ombitasvir/paritaprevir/ritonavir tablets must be administered with dasabuvir tablets.

Recommended Dose in Adults

The recommended oral dose of VIEKIRA PAK is two paritaprevir/ritonavir/ombitasvir 75/50/12.5 mg tablets once daily (in the morning) and one dasabuvir 250 mg tablet twice daily (morning and evening).

VIEKIRA PAK is used in non-cirrhotic genotype 1b-infected patients, and can be considered as a therapeutic option for non-cirrhotic, treatment-naïve patients with genotype 1a infection (see CLINICAL TRIALS and Table 15).

Patients should be instructed to swallow the tablets whole (i.e. patients should not chew, break or dissolve the tablets).

To maximise absorption, VIEKIRA PAK should be taken with food without regard to fat or calorie content (see PHARMACOLOGY).

Table 15 shows the recommended treatment regimen and duration based on patient population.

Table 15: Treatment Regimen and Duration by Patient Population

Patient Population	Treatment	Duration	Ribavirin Dosage
Genotype 1b, without cirrhosis	VIEKIRA PAK	12 weeks	NA
Genotype 1a, without cirrhosis	VIEKIRA PAK- RBV*	12 weeks	< 75 kg = 1000 mg ≥ 75 kg = 1200 mg Ribavirin is to be taken in two doses, morning and evening
Genotype 1 with cirrhosis	VIEKIRA PAK- RBV	12 weeks [†]	< 75 kg = 1000 mg ≥ 75 kg = 1200 mg Ribavirin is to be taken in two doses, morning and evening
<p>* VIEKIRA PAK without ribavirin can be considered as a therapeutic option for treatment-naïve patients with genotype 1a infection without cirrhosis (see CLINICAL TRIALS). Treatment decision should be guided by an assessment of the potential benefits and risks and available alternative therapies for the individual patient.</p> <p>† 24 weeks of VIEKIRA PAK–RBV is recommended for patients with genotype 1a-infection with cirrhosis who have had a previous null response to pegIFN and ribavirin (see CLINICAL TRIALS). Note: VIEKIRA PAK-RBV is recommended in patients with an unknown genotype 1 subtype or with mixed genotype 1 infection.</p>			

VIEKIRA PAK should be taken as directed for the prescribed duration, without interruption.

Missed Dose

Inform patients that in case a dose of paritaprevir, ritonavir, ombitasvir is missed, the prescribed dose can be taken within 12 hours.

In case a dose of dasabuvir is missed, the prescribed dose can be taken within 6 hours.

If more than 12 hours has passed since ombitasvir, paritaprevir, ritonavir is usually taken or more than 6 hours has passed since dasabuvir is usually taken, the missed dose should NOT be taken and the patient should take the next dose as per the usual dosing schedule.

Instruct patients not to take more than their prescribed dose of VIEKIRA PAK to make up for a missed dose.

Use in Special Populations

Hepatic Impairment

No dose adjustment of VIEKIRA PAK is required in patients with mild hepatic impairment (Child-Pugh A). The safety and efficacy of VIEKIRA PAK have not been established in HCV-infected patients with moderate hepatic impairment (Child-Pugh B); however, no dose adjustment is expected to be required based on pharmacokinetic studies. VIEKIRA PAK is contraindicated in patients with severe hepatic impairment (Child-Pugh C).

Renal Impairment

Based on the pharmacokinetic data in HCV uninfected subjects (n=24), no dose adjustment of VIEKIRA PAK is recommended in subjects with mild, moderate or severe renal impairment. The efficacy and safety of VIEKIRA PAK have not been evaluated in HCV-infected subjects with moderate or severe renal impairment. VIEKIRA PAK has not been studied in patients on dialysis.

OVERDOSAGE

The highest documented single dose administered to healthy volunteers was 400 mg for paritaprevir (with 100 mg ritonavir), 200 mg for ritonavir (with 100 mg paritaprevir), 350 mg for ombitasvir and 2000 mg for dasabuvir. In case of overdose, it is recommended that the patient be monitored for any signs or symptoms of adverse reactions and appropriate symptomatic treatment instituted immediately.

For information on the management of overdose in Australia contact the Poison Information Centre on 131126 and in New Zealand call 0800 764 766.

PRESENTATION AND STORAGE CONDITIONS

Presentation

Paritaprevir/ritonavir/ombitasvir 75/50/12.5 mg tablets are pink-coloured, film-coated, oblong biconvex shaped, debossed with "AV1" on one side.

Dasabuvir 250 mg tablets are beige-coloured, film-coated, oval-shaped, debossed with "AV2" on one side

VIEKIRA PAK is dispensed in a monthly carton for a total of 28 days of therapy. Each monthly carton contains four weekly cartons. Each weekly carton contains seven daily dose packs. Each daily dose pack contains four tablets: two 75 mg/50 mg/12.5 mg paritaprevir/ritonavir/ombitasvir tablets and two 250mg dasabuvir tablets in PVC/PE/PCTFE(Aclar)/Al blisters, and indicates which tablets need to be taken in the morning and evening.

Attachment 1: Product information AusPAR VIEKIRA PAK - Paritaprevir / Ritonavir / Ombitasvir and Dasabuvir (as Sodium Salt) - AbbVie Pty Ltd - PM-2014-01436-1-2 FINAL 24 March 2017. This Product Information was approved at the time this AusPAR was published.

Storage Conditions

Store below 25°C in a dry place.

NAME AND ADDRESS OF THE SPONSOR

AbbVie Pty Ltd
Level 7, 241 O’Riordan Street
Mascot NSW 2020
Australia

AbbVie Limited
6th Floor, 156-158 Victoria St
Wellington, 6011
New Zealand

POISON SCHEDULE OF THE MEDICINE

Prescription Only Medicine

Schedule 4

DATE OF FIRST INCLUSION IN THE ARTG

10 July 2015

DATE OF MOST RECENT AMENDMENT

N/A